

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
4 April 2002 (04.04.2002)

PCT

(10) International Publication Number  
**WO 02/26734 A1**

(51) International Patent Classification<sup>7</sup>: **C07D 409/12**,  
409/14, 241/08, 403/10, 401/14, 491/113, 401/12, 405/14,  
A61K 31/495, A61P 7/02 // (C07D 491/113, 221:00,  
317:00)

94404 (US). **SCARBOROUGH, Robert, M.** [US/US];  
22 Greenbrier Court, Half Moon Bay, CA 94019 (US).

(21) International Application Number: PCT/US01/30313

(22) International Filing Date: 1 October 2001 (01.10.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/236,393 29 September 2000 (29.09.2000) US

(71) Applicant (for all designated States except US): **COR  
THERAPEUTICS, INC.** [US/US]; 256 East Grand  
Avenue, South San Francisco, CA 94080 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ZHU, Bing-Yan**  
[CA/US]; 135 Lois Lane, Palo Alto, CA 94303 (US).  
**SU, Ting** [US/US]; 3222 Upper Lock Avenue, Belmont,  
CA 94002 (US). **LI, Wenhao** [CN/US]; P.O. Box 1993,  
South San Francisco, CA 94083 (US). **GOLDMAN,  
Erick, A.** [US/US]; 1520 Francisco Street, Berkeley, CA  
94702 (US). **ZHANG, Penglie** [CN/US]; 251 Winchester  
Court, Foster City, CA 94404 (US). **JIA, Zhaozhong,  
Jon** [CN/US]; 1576-E Marina Court, San Mateo, CA

(74) Agents: **LEE, Christine, S.** et al.; Morgan, Lewis & Bock-  
ius LLP, 1800 M. Street, NW, Washington, DC 20036 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,  
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU,  
ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,  
TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: PIPERAZIN-2-ONE AMIDES AS INHIBITORS OF FACTOR Xa

(57) Abstract: Novel piperazin-2-one containing compounds of general formulae (I) or (II), including their pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivative having activity against mammalian factor Xa are described. Compositions containing such compounds are also described. The compounds and the compositions are useful *in vitro* or *in vivo* for preventing or treating conditions in mammals characterized by undesired thrombosis.

WO 02/26734 A1

## PIPERAZIN-2-ONE AMIDES AS INHIBITORS OF FACTOR XA

Field of the Invention

The invention relates to novel piperazin-2-one-containing compounds including  
5 their pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug  
derivatives, and pharmaceutically acceptable compositions thereof which are potent and  
highly selective inhibitors of isolated factor Xa or when assembled in the  
prothrombinase complex. These compounds show selectivity for factor Xa versus other  
proteases of the coagulation (e.g. thrombin, fVIIa, fIXa) or the fibrinolytic cascades  
10 (e.g. plasminogen activators, plasmin). In another aspect, the present invention relates  
to novel piperazin-2-one-containing compounds including their pharmaceutically  
acceptable isomers, salts, hydrates, solvates and prodrug derivatives factor Xa-  
inhibiting compounds, and pharmaceutically acceptable compositions thereof which are  
useful as potent and specific inhibitors of blood coagulation in mammals. In yet  
15 another aspect, the invention relates to methods for using these inhibitors as therapeutic  
agents for disease states in mammals characterized by undesired thrombosis or  
coagulation disorders.

Background of the Invention

20 Hemostasis, the control of bleeding, occurs by surgical means, or by the  
physiological properties of vasoconstriction and coagulation. The invention is  
particularly concerned with blood coagulation and ways in which it assists in  
maintaining the integrity of mammalian circulation after injury, inflammation, disease,  
congenital defect, dysfunction or other disruption. Although platelets and blood  
25 coagulation are both involved in thrombus formation, certain components of the  
coagulation cascade are primarily responsible for the amplification or acceleration of  
the processes involved in platelet aggregation and fibrin deposition.

Thrombin is a key enzyme in the coagulation cascade as well as in hemostasis.  
Thrombin plays a central role in thrombosis through its ability to catalyze the  
30 conversion of fibrinogen into fibrin and through its potent platelet activation activity.  
Direct or indirect inhibition of thrombin activity has been the focus of a variety of

recent anticoagulant strategies as reviewed by Claeson, G., "Synthetic Peptides and Peptidomimetics as Substrates and Inhibitors of Thrombin and Other Proteases in the Blood Coagulation System", *Blood Coag. Fibrinol.* 5, 411-436 (1994). Several classes of anticoagulants currently used in the clinic directly or indirectly affect thrombin (i.e. 5 heparins, low-molecular weight heparins, heparin-like compounds and coumarins).

A prothrombinase complex, including Factor Xa (a serine protease, the activated form of its Factor X precursor and a member of the calcium ion binding, gamma carboxyglutamyl (Gla)-containing, vitamin K dependent, blood coagulation glycoprotein family), converts the zymogen prothrombin into the active procoagulant 10 thrombin. Unlike thrombin, which acts on a variety of protein substrates as well as at a specific receptor, factor Xa appears to have a single physiologic substrate, namely prothrombin. Since one molecule of factor Xa may be able to generate up to 138 molecules of thrombin (Elodi et al., *Thromb. Res.* 15, 617-619 (1979)), direct inhibition of factor Xa as a way of indirectly inhibiting the formation of thrombin may be an 15 efficient anticoagulant strategy. Therefore, it has been suggested that compounds which selectively inhibit factor Xa may be useful as *in vitro* diagnostic agents, or for therapeutic administration in certain thrombotic disorders, see e.g., WO 94/13693.

Polypeptides derived from hematophagous organisms have been reported which are highly potent and specific inhibitors of factor Xa. United States Patent 4,588,587 20 describes anticoagulant activity in the saliva of the Mexican leech, *Haementeria officinalis*. A principal component of this saliva was shown to be the polypeptide factor Xa inhibitor, antistasin (ATS), by Nutt, E. et al., "The Amino Acid Sequence of Antistasin, a Potent Inhibitor of Factor Xa Reveals a Repeated Internal Structure", *J. Biol. Chem.*, 263, 10162-10167 (1988). Another potent and highly specific inhibitor of 25 Factor Xa, called tick anticoagulant peptide (TAP), has been isolated from the whole body extract of the soft tick *Ornithodoros moubata*, as reported by Waxman, L., et al., "Tick Anticoagulant Peptide (TAP) is a Novel Inhibitor of Blood Coagulation Factor Xa" *Science*, 248, 593-596 (1990).

Factor Xa inhibitory compounds which are not large polypeptide-type inhibitors 30 have also been reported including: Tidwell, R.R. et al., "Strategies for Anticoagulation With Synthetic Protease Inhibitors. Xa Inhibitors Versus Thrombin Inhibitors",

- Thromb. Res., 19, 339-349 (1980); Turner, A.D. *et al.*, "p-Amidino Esters as Irreversible Inhibitors of Factor IXa and Xa and Thrombin", Biochemistry, 25, 4929-4935 (1986); Hitomi, Y. *et al.*, "Inhibitory Effect of New Synthetic Protease Inhibitor (FUT-175) on the Coagulation System", Haemostasis, 15, 164-168 (1985);
- 5 Sturzebecher, J. *et al.*, "Synthetic Inhibitors of Bovine Factor Xa and Thrombin. Comparison of Their Anticoagulant Efficiency", Thromb. Res., 54, 245-252 (1989); Kam, C.M. *et al.*, "Mechanism Based Isocoumarin Inhibitors for Trypsin and Blood Coagulation Serine Proteases: New Anticoagulants", Biochemistry, 27, 2547-2557 (1988); Hauptmann, J. *et al.*, "Comparison of the Anticoagulant and Antithrombotic
- 10 Effects of Synthetic Thrombin and Factor Xa Inhibitors", Thromb. Haemost., 63, 220-223 (1990); and the like.

Others have reported Factor Xa inhibitors which are small molecule organic compounds, such as nitrogen containing heterocyclic compounds which have amidino substituent groups, wherein two functional groups of the compounds can bind to Factor

15 Xa at two of its active sites. For example, WO 98/28269 describes pyrazole compounds having a terminal C(=NH)-NH<sub>2</sub> group; WO 97/21437 describes benzimidazole compounds substituted by a basic radical which are connected to a naphthyl group via a straight or branched chain alkylene, -C(=O) or -S(=O)<sub>2</sub> bridging group; WO 99/10316 describes compounds having a 4-phenyl-N-alkylamidino-

20 piperidine and 4-phenoxy-N-alkylamidino-piperidine group connected to a 3-amidinophenyl group via a carboxamidealkyleneamino bridge; and EP 798295 describes compounds having a 4-phenoxy-N-alkylamidino-piperidine group connected to an amidinonaphthyl group via a substituted or unsubstituted sulfonamide or carboxamide bridging group.

25 There exists a need for effective therapeutic agents for the regulation of hemostasis, and for the prevention and treatment of thrombus formation and other pathological processes in the vasculature induced by thrombin such as restenosis and inflammation. In particular, there continues to be a need for compounds which selectively inhibit factor Xa or its precursors. Compounds are needed which selectively


30 or preferentially bind to Factor Xa. Compounds with a higher affinity for binding to

Factor Xa than to thrombin are desired, especially those compounds having good bioavailability or other pharmacologically desirable properties.

## Summary of the Invention


5 The present invention relates to novel piperazin-2-one-containing compounds including their pharmaceutically acceptable isomers, salts, hydrates, solvate and prodrug derivatives, which have particular biological properties and are useful as potent and specific inhibitors of blood coagulation in mammals. According to the invention, the compounds can act as potent and highly selective inhibitors of isolated Factor Xa or  
10 when assembled in the prothrombinase complex. The invention also provides compositions containing such compounds. The compounds of the invention may be used as diagnostic reagents or as therapeutic reagents for disease states in mammals which have coagulation disorders. Thus, the invention further provides methods for preventing or treating a condition in a mammal characterized by undesired thrombosis  
15 by administration of a therapeutically effective amount of a compound of the invention and a pharmaceutically acceptable carrier. Optionally, the methods of the invention comprise administering a pharmaceutical composition of the invention in combination with an additional therapeutic agent such as an antithrombotic and/or a thrombolytic agent and/or an anticoagulant. According to the invention, such conditions include, for  
20 example, any thrombotically mediated acute coronary or cerebrovascular syndrome, any thrombotic syndrome occurring in the venous system, any coagulopathy, and any thrombotic complications associated with extracorporeal circulation or instrumentation, and for the inhibition of coagulation in biological samples (e.g. stored blood products and samples).

25        The invention provides a compound of the general formulae I or II:



wherein:

A is a member selected from the group consisting of:




- $R^{1a}$ ,  $R^{1b}$ ,  $R^{1d}$ , and  $R^{1e}$  are each independently a H,  $C_{1-6}$  alkyl,  $C_{2-6}$  alkenyl,  $C_{2-6}$  alkynyl,  $C_{3-8}$  cycloalkyl, aryl,  $-C_{1-6}$  alkylaryl, heterocyclyl,  $-C_{1-6}$  alkylheterocyclyl,  $-(CH_2)_{1-6}OH$ ,  $-(CH_2)_{1-6}OC_{1-6}$  alkyl,  $-(CH_2)_{1-6}NH_2$ ,  $-(CH_2)_{1-6}NHC_{1-6}$  alkyl,  $-(CH_2)_{1-6}N(C_{1-6} \text{ alkyl})_2$ ,  $-(CH_2)_{1-6}CHNH(COOH)$ ,  $-(CH_2)_{1-6}NHC(=O)C_{1-6}$  alkyl,  $-(CH_2)_{1-6}CHO$ ,  $-(CH_2)_{1-6}C(=O)OH$ ,  $-(CH_2)_{1-6}C(=O)OC_{1-6}$  alkyl, or  $-(CH_2)_{1-6}C(=O)NH_2$ ; wherein  $R^{1a}$ ,  $R^{1b}$ ,  $R^{1d}$ , or  $R^{1e}$  is optionally substituted with at least one of halo, alkyl, alkylideneamine, arylideneamine, cyano, hydroxy, alkoxy, amino, amidino, guanidino, imino, amido, acid, ester, keto, aldehyde, dioxolane, furanyl, piperidiny, piperaziny, pyrrolidiny, aryl, morpholiny, and thiomorpholinyldioxide; or  $R^{1a}$  and  $R^{1b}$  or  $R^{1a}$  and  $R^{1c}$  or  $R^{1a}$  and  $R^{1d}$  or  $R^{1d}$  and  $R^{1e}$  taken together with the nitrogen atom to which they are each attached can form a substituted or unsubstituted 3 to 8 membered heterocyclic or heteroaromatic amine group which, optionally, contains at least one other heteroatom of N, O or S;
- wherein  $R^{1a}$ ,  $R^{1b}$ ,  $R^{1d}$ , or  $R^{1e}$  is optionally substituted with at least one of halo, alkyl, alkylideneamine, arylideneamine, cyano, hydroxy, alkoxy, amino, amidino, guanidino, imino, amido, acid, ester, keto, aldehyde, dioxolane, furanyl, piperidiny, piperaziny, pyrrolidiny, aryl, morpholiny, and thiomorpholinyldioxide;
- $R^{1c}$  is H,  $C_{1-6}$  alkyl or  $C_{3-8}$  cycloalkyl;

$R^{2a}$ ,  $R^{2b}$  and  $R^{2c}$  are each independently a H,  $C_{1-6}$  alkyl,  $C_{2-6}$  alkenyl,  $C_{2-6}$  alkynyl,  $C_{3-8}$  cycloalkyl, aryl,  $-C_{1-6}$  alkylaryl, heterocyclyl,  $-C_{1-6}$  alkylheterocyclyl,  $-(CH_2)_{1-6}OH$ , -

- (CH<sub>2</sub>)<sub>1-6</sub>OC<sub>1-6</sub> alkyl, -(CH<sub>2</sub>)<sub>1-6</sub>NH<sub>2</sub>, -(CH<sub>2</sub>)<sub>1-6</sub>NHC<sub>1-6</sub> alkyl, -(CH<sub>2</sub>)<sub>1-6</sub>N(C<sub>1-6</sub> alkyl)<sub>2</sub>, -(CH<sub>2</sub>)<sub>1-6</sub>CHNH(COOH), -(CH<sub>2</sub>)<sub>1-6</sub>NHC(=O)C<sub>1-6</sub> alkyl, -(CH<sub>2</sub>)<sub>1-6</sub>CHO, -(CH<sub>2</sub>)<sub>1-6</sub>C(=O)OH, -(CH<sub>2</sub>)<sub>1-6</sub>C(=O)OC<sub>1-6</sub>alkyl, or -(CH<sub>2</sub>)<sub>1-6</sub>C(=O)NH<sub>2</sub>; wherein R<sup>2a</sup>, R<sup>2b</sup> and R<sup>2c</sup> is optionally substituted with at least one of halo, alkyl, alkylideneamine, arylidenamine, cyano, hydroxy, alkoxy, amino, amidino, guanidino, imino, amido, acid, ester, keto, aldehyde, dioxolane, furanyl, piperidinyl, piperazinyl, pyrrolidinyl, aryl, morpholinyl, and thiomorpholinyl dioxide; or R<sup>2a</sup> and R<sup>2b</sup> or R<sup>1a</sup>, as set forth above, and R<sup>2a</sup> or R<sup>1a</sup>, as set forth above, and R<sup>2b</sup> taken together with the nitrogen atom to which they are each attached can form a substituted or unsubstituted 3 to 8 membered heterocyclic or heteroaromatic amine group which, optionally, contains at least one other heteroatom of N, O or S; wherein R<sup>2a</sup>, R<sup>2b</sup> or R<sup>2c</sup> is optionally substituted with at least one of halo, alkyl, alkylideneamine, arylidenamine, cyano, hydroxy, alkoxy, amino, amidino, guanidino, imino, amido, acid, ester, keto, aldehyde, dioxolane, furanyl, piperidinyl, piperazinyl, pyrrolidinyl, aryl, morpholinyl, and thiomorpholinyl dioxide;

- R is, in each occurrence, independently, H, -C<sub>1-6</sub>alkyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-6</sub>alkyl-OC<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkyl-O(CH<sub>2</sub>)<sub>1-4</sub>-COOH, -C<sub>0-6</sub>alkyl-O(CH<sub>2</sub>)<sub>1-4</sub>-C(=O)OC<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkylCOOH, -C<sub>0-6</sub>alkylCO<sub>2</sub>C<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkylOC<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkylOH, -C<sub>0-6</sub>alkylCONH<sub>2</sub>, -C<sub>0-6</sub>alkylCONHC<sub>0-6</sub>alkyl, -C<sub>0-6</sub>alkylCON(C<sub>0-6</sub>alkyl)<sub>2</sub>, -C<sub>0-6</sub>alkylCON(CH<sub>2</sub>)<sub>2-6</sub>, -C<sub>0-6</sub>alkylCON(CH<sub>2</sub>CH<sub>2</sub>)<sub>2</sub>O, -C<sub>0-6</sub>alkylCON(CH<sub>2</sub>CH<sub>2</sub>)<sub>2</sub>SO<sub>2</sub>, -C<sub>0-6</sub>alkylCONHaryl, -C<sub>0-6</sub>alkylNH<sub>2</sub>, -C<sub>0-6</sub>alkylNH(C<sub>1-6</sub>alkyl) or -C<sub>0-6</sub>alkylN(C<sub>1-6</sub>alkyl)<sub>2</sub>.


- Q is a member selected from the group consisting of:



$Y$  is S;

5  $R^1$  is H, -Cl, -Br, -I, -F, -OCF<sub>3</sub>, alkyl, hydroxy, alkoxy, amino, thiol, thioalkyl, thioaryl, or piperiziny;

$J^1$  is a member selected from the group consisting of:



10  $X$  is O or S;


$R^2$  is H, -Cl, -Br, -I, -F or -OC<sub>1-6</sub>alkyl;

$R^3$  is H, -Cl, -Br, -I, -F, -OC<sub>1-6</sub>alkyl, -NHC<sub>1-6</sub>acyl, -NO<sub>2</sub>, -NHSO<sub>2</sub>C<sub>1-4</sub>alkyl, -CN, -NH<sub>2</sub>, -CONH<sub>2</sub>, -SO<sub>2</sub>C<sub>1-6</sub>alkyl, -SO<sub>2</sub>NH<sub>2</sub>, -CO<sub>2</sub>C<sub>1-6</sub>alkyl or -O(CH<sub>2</sub>)<sub>1-4</sub>COOH;

15

$R^4$  and  $R^5$  are each independently H, -Cl, -Br, -I, -F or -OC<sub>1-6</sub>alkyl;

$J^2$  is a member selected from the group consisting of:



20

$Z$  is -NR<sup>6</sup>-, -O- or -S-;



R<sup>6</sup> is H, C<sub>1-6</sub>alkyl or C<sub>3-8</sub>cycloalkyl;

R<sup>7</sup> and R<sup>8</sup> are independently H, -Cl, -Br, -I or -F, where at least one of R<sup>7</sup> and R<sup>8</sup> is not  
5 hydrogen; and

R<sup>9</sup> and R<sup>10</sup> are independently H, -Cl, -Br, -I or -F, where at least one of R<sup>9</sup> and R<sup>10</sup> is  
not hydrogen;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and  
10 prodrug derivatives thereof.

### Detailed Description of the Invention

#### Definitions

In accordance with the present invention and as used herein, the following terms  
15 are defined with the following meanings, unless explicitly stated otherwise.

The term "alkenyl" refers to a trivalent straight chain or branched chain  
unsaturated aliphatic radical. The term "alkynyl" (or "alkinyl") refers to a straight or  
branched chain aliphatic radical that includes at least two carbons joined by a triple  
bond. If no number of carbons is specified alkenyl and alkynyl each refer to radicals  
20 having from 2-12 carbon atoms.

The term "alkyl" refers to saturated aliphatic groups including straight-chain,  
branched-chain and cyclic groups having the number of carbon atoms specified, or if no  
number is specified, having up to 12 carbon atoms. The term "cycloalkyl" as used  
herein refers to a mono-, bi-, or tricyclic aliphatic ring having 3 to 14 carbon atoms and  
25 preferably 3 to 7 carbon atoms.

As used herein, the terms "carbocyclic ring structure" and "C<sub>3-16</sub> carbocyclic  
mono, bicyclic or tricyclic ring structure" or the like are each intended to mean stable  
ring structures having only carbon atoms as ring atoms wherein the ring structure is a  
substituted or unsubstituted member selected from the group consisting of: a stable  
30 monocyclic ring which is aromatic ring ("aryl") having six ring atoms; a stable  
monocyclic non-aromatic ring having from 3 to 7 ring atoms in the ring; a stable

bicyclic ring structure having a total of from 7 to 12 ring atoms in the two rings wherein the bicyclic ring structure is selected from the group consisting of ring structures in which both of the rings are aromatic, ring structures in which one of the rings is aromatic and ring structures in which both of the rings are non-aromatic; and a  
5 stable tricyclic ring structure having a total of from 10 to 16 atoms in the three rings wherein the tricyclic ring structure is selected from the group consisting of: ring structures in which three of the rings are aromatic, ring structures in which two of the rings are aromatic and ring structures in which three of the rings are non-aromatic. In each case, the non-aromatic rings when present in the monocyclic, bicyclic or tricyclic  
10 ring structure may independently be saturated, partially saturated or fully saturated. Examples of such carbocyclic ring structures include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, adamantyl, cyclooctyl, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane (decalin), 2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, or  
15 tetrahydronaphthyl (tetralin). Moreover, the ring structures described herein may be attached to one or more indicated pendant groups via any carbon atom which results in a stable structure. The term "substituted" as used in conjunction with carbocyclic ring structures means that hydrogen atoms attached to the ring carbon atoms of ring structures described herein may be substituted by one or more of the substituents  
20 indicated for that structure if such substitution(s) would result in a stable compound.

The term "aryl" which is included with the term "carbocyclic ring structure" refers to an unsubstituted or substituted aromatic ring, substituted with one, two or three substituents selected from loweralkoxy, loweralkyl, loweralkylamino, hydroxy, halogen, cyano, hydroxyl, mercapto, nitro, thioalkoxy, carboxaldehyde, carboxyl,  
25 carboalkoxy and carboxamide, including but not limited to carbocyclic aryl, heterocyclic aryl, and biaryl groups and the like, all of which may be optionally substituted. Examples of suitable aryl groups include, but are not limited to, phenyl, pyridyl, thiophenyl, halophenyl, loweralkylphenyl, naphthyl, biphenyl, phenanthrenyl and naphthacenyl.

30 The term "arylalkyl" which is included with the term "carbocyclic aryl" refers to one, two, or three aryl groups having the number of carbon atoms designated, appended

to an alkyl group having the number of carbon atoms designated. Suitable arylalkyl groups include, but are not limited to, benzyl, picolyl, naphthylmethyl, phenethyl, benzyhydril, trityl, and the like, all of which may be optionally substituted.

As used herein, the term "heterocyclic ring" or "heterocyclic ring system" is  
5 intended to mean a substituted or unsubstituted member selected from the group consisting of stable monocyclic ring having from 5-7 members in the ring itself and having from 1 to 4 hetero ring atoms selected from the group consisting of N, O and S; a stable bicyclic ring structure having a total of from 7 to 12 atoms in the two rings wherein at least one of the two rings has from 1 to 4 hetero atoms selected from N, O  
10 and S, including bicyclic ring structures wherein any of the described stable monocyclic heterocyclic rings is fused to a hexane or benzene ring; and a stable tricyclic heterocyclic ring structure having a total of from 10 to 16 atoms in the three rings wherein at least one of the three rings has from 1 to 4 hetero atoms selected from the group consisting of N, O and S. Any nitrogen and sulfur atoms present in a  
15 heterocyclic ring of such a heterocyclic ring structure may be oxidized. Unless indicated otherwise the terms "heterocyclic ring" or "heterocyclic ring system" include aromatic rings, as well as non-aromatic rings which can be saturated, partially saturated or fully saturated non-aromatic rings. Also, unless indicated otherwise the term "heterocyclic ring system" includes ring structures wherein all of the rings contain at  
20 least one hetero atom as well as structures having less than all of the rings in the ring structure containing at least one hetero atom, for example bicyclic ring structures wherein one ring is a benzene ring and one of the rings has one or more hetero atoms are included within the term "heterocyclic ring systems" as well as bicyclic ring structures wherein each of the two rings has at least one hetero atom. Moreover, the  
25 ring structures described herein may be attached to one or more indicated pendant groups via any hetero atom or carbon atom which results in a stable structure. Further, the term "substituted" means that one or more of the hydrogen atoms on the ring carbon atom(s) or nitrogen atom(s) of the each of the rings in the ring structures described herein may be replaced by one or more of the indicated substituents if such  
30 replacement(s) would result in a stable compound. Nitrogen atoms in a ring structure may be quaternized, but such compounds are specifically indicated or are included

within the term "a pharmaceutically acceptable salt" for a particular compound. When the total number of O and S atoms in a single heterocyclic ring is greater than 1, it is preferred that such atoms not be adjacent to one another. Preferably, there are no more than 1 O or S ring atoms in the same ring of a given heterocyclic ring structure.

5        Examples of monocyclic and bicyclic heterocyclic ring systems, in alphabetical order, are acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazalinyl, carbazolyl, 4aH-carbazolyl, carbolinyl, chromanyl, chromenyl, cinnolinyl, decahydroquinolinyl, 2H,6H-1,5,2-  
10 dithiazinyl, dihydrofuro[2,3-b]tetrahydrofuran, furanyl, furazanyl, imidazolidinyl, imidazolyl, imidazolyl, 1H-indazolyl, indolinyl, indoliziny, indolyl, 3H-indolyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindolinyl, isoindolyl, isoquinolinyl, (benzimidazolyl), isothiazolyl, isoxazolyl, morpholinyl, naphthyridinyl, octahydroisoquinolinyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl,  
15 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl, oxazolyl, oxazolidinyl, pyrimidinyl, phenanthridinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, phenoxathiinyl, phenoxazinyl, phthalazinyl, piperazinyl, piperidinyl, pteridinyl, purinyl, pyranyl, pyrazinyl, pyroazolidinyl, pyrazolinyl, pyrazolyl, pyridazinyl, pyridooxazole, pyridoimidazole, pyridothiazole, pyridinyl, pyridyl, pyrimidinyl, pyrrolidinyl,  
20 pyrrolinyl, 2H-pyrrolyl, pyrrolyl, quinazolinyl, quinolinyl, 4H-quinoliziny, quinoxalinyl, quinuclidinyl, tetrahydrofuranyl, tetrahydroisoquinolinyl, tetrahydroquinolinyl, 6H-1,2,5-thiadiazinyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, thianthrenyl, thiazolyl, thienyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl, thiophenyl, triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl,  
25 1,2,5-triazolyl, 1,3,4-triazolyl and xanthenyl. Preferred heterocyclic ring structures include, but are not limited to, pyridinyl, furanyl, thienyl, pyrrolyl, pyrazolyl, pyrrolidinyl, imidazolyl, indolyl, benzimidazolyl, 1H-indazolyl, oxazolyl, or isatinoyl. Also included are fused ring and spiro compounds containing, for example, the above heterocyclic ring structures.

30        As used herein the term "aromatic heterocyclic ring system" has essentially the same definition as for the monocyclic and bicyclic ring systems except that at least one

ring of the ring system is an aromatic heterocyclic ring or the bicyclic ring has an aromatic or non-aromatic heterocyclic ring fused to an aromatic carbocyclic ring structure.

The terms "halo" or "halogen" as used herein refer to Cl, Br, F or I substituents.

- 5 The term "haloalkyl", and the like, refer to an aliphatic carbon radicals having at least one hydrogen atom replaced by a Cl, Br, F or I atom, including mixtures of different halo atoms. Trihaloalkyl includes trifluoromethyl and the like as preferred radicals, for example.

The term "methylene" refers to  $-\text{CH}_2-$ .

- 10 The term "pharmaceutically acceptable salts" includes salts of compounds derived from the combination of a compound and an organic or inorganic acid. These compounds are useful in both free base and salt form. In practice, the use of the salt form amounts to use of the base form; both acid and base addition salts are within the scope of the present invention.

- 15 "Pharmaceutically acceptable acid addition salt" refers to salts retaining the biological effectiveness and properties of the free bases and which are not biologically or otherwise undesirable, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic  
20 acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like.

- "Pharmaceutically acceptable base addition salts" include those derived from inorganic bases such as sodium, potassium, lithium, ammonium, calcium, magnesium,  
25 iron, zinc, copper, manganese, aluminum salts and the like. Particularly preferred are the ammonium, potassium, sodium, calcium and magnesium salts. Salts derived from pharmaceutically acceptable organic nontoxic bases include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as  
30 isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-diethylaminoethanol, trimethamine, dicyclohexylamine, lysine,


arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, methylglucamine, theobromine, purines, piperizine, piperidine, N-ethylpiperidine, polyamine resins and the like. Particularly preferred organic nontoxic bases are isopropylamine, diethylamine, ethanolamine, trimethamine,  
 5 dicyclohexylamine, choline, and caffeine.

"Biological property" for the purposes herein means an *in vivo* effector or antigenic function or activity that is directly or indirectly performed by a compound of the invention that are often shown by *in vitro* assays. Effector functions include receptor or ligand binding, any enzyme activity or enzyme modulatory activity, any  
 10 carrier binding activity, any hormonal activity, any activity in promoting or inhibiting adhesion of cells to an extracellular matrix or cell surface molecules, or any structural role. Antigenic functions include possession of an epitope or antigenic site that is capable of reacting with antibodies raised against it.

In the compounds of the invention, carbon atoms bonded to four non-identical  
 15 substituents are asymmetric. Accordingly, the compounds may exist as diastereoisomers, enantiomers or mixtures thereof. The syntheses described herein may employ racemates, enantiomers or diastereomers as starting materials or intermediates. Diastereomeric products resulting from such syntheses may be separated by chromatographic or crystallization methods, or by other methods known in the art.  
 20 Likewise, enantiomeric product mixtures may be separated using the same techniques or by other methods known in the art. Each of the asymmetric carbon atoms, when present in the compounds of the invention, may be in one of two configurations (R or S) and both are within the scope of the present invention.


## 25 Compounds

The invention provides a compound of the general formulae I or II:



wherein:

A is a member selected from the group consisting of:



- $R^{1a}$ ,  $R^{1b}$ ,  $R^{1d}$ , and  $R^{1e}$  are each independently a H,  $C_{1-6}$  alkyl,  $C_{2-6}$  alkenyl,  $C_{2-6}$  alkynyl,  $C_{3-8}$  cycloalkyl, aryl,  $-C_{1-6}$  alkylaryl, heterocyclyl,  $-C_{1-6}$  alkylheterocyclyl,  $-(CH_2)_{1-6}OH$ ,  $-(CH_2)_{1-6}OC_{1-6}$  alkyl,  $-(CH_2)_{1-6}NH_2$ ,  $-(CH_2)_{1-6}NHC_{1-6}$  alkyl,  $-(CH_2)_{1-6}N(C_{1-6} \text{ alkyl})_2$ ,  $-(CH_2)_{1-6}CHNH(COOH)$ ,  $-(CH_2)_{1-6}NHC(=O)C_{1-6}$  alkyl,  $-(CH_2)_{1-6}CHO$ ,  $-(CH_2)_{1-6}C(=O)OH$ ,  $-(CH_2)_{1-6}C(=O)OC_{1-6}$  alkyl, or  $-(CH_2)_{1-6}C(=O)NH_2$ ; wherein  $R^{1a}$ ,  $R^{1b}$ ,  $R^{1d}$ , or  $R^{1e}$  is optionally substituted with at least one of halo, alkyl, alkylideneamine, arylidenamine, cyano, hydroxy, alkoxy, amino, amidino, guanidino, imino, amido, acid, ester, keto, aldehyde, dioxolane, furanyl, piperidiny, piperaziny, pyrrolidiny, aryl, morpholiny, and thiomorpholinyldioxide; or  $R^{1a}$  and  $R^{1b}$  or  $R^{1a}$  and  $R^{1c}$  or  $R^{1a}$  and  $R^{1d}$  or  $R^{1d}$  and  $R^{1e}$  taken together with the nitrogen atom to which they are each attached can form a substituted or unsubstituted 3 to 8 membered heterocyclic or heteroaromatic amine group which, optionally, contains at least one other heteroatom of N, O or S; wherein  $R^{1a}$ ,  $R^{1b}$ ,  $R^{1d}$ , or  $R^{1e}$  is optionally substituted with at least one of halo, alkyl, alkylideneamine, arylidenamine, cyano, hydroxy, alkoxy, amino, amidino, guanidino, imino, amido, acid, ester, keto, aldehyde, dioxolane, furanyl, piperidiny, piperaziny, pyrrolidiny, aryl, morpholiny, and thiomorpholinyldioxide;
- $R^{1c}$  is H,  $C_{1-6}$  alkyl or  $C_{3-8}$  cycloalkyl;

$R^{2a}$ ,  $R^{2b}$  and  $R^{2c}$  are each independently a H,  $C_{1-6}$  alkyl,  $C_{2-6}$  alkenyl,  $C_{2-6}$  alkynyl,  $C_{3-8}$  cycloalkyl, aryl,  $-C_{1-6}$  alkylaryl, heterocyclyl,  $-C_{1-6}$  alkylheterocyclyl,  $-(CH_2)_{1-6}OH$ ,  $-(CH_2)_{1-6}OC_{1-6}$  alkyl,  $-(CH_2)_{1-6}NH_2$ ,  $-(CH_2)_{1-6}NHC_{1-6}$  alkyl,  $-(CH_2)_{1-6}N(C_{1-6} \text{ alkyl})_2$ ,  $-(CH_2)_{1-6}CHNH(COOH)$ ,  $-(CH_2)_{1-6}NHC(=O)C_{1-6}$  alkyl,  $-(CH_2)_{1-6}CHO$ ,  $-(CH_2)_{1-6}C(=O)OH$ ,  $-(CH_2)_{1-6}C(=O)OC_{1-6}$  alkyl, or  $-(CH_2)_{1-6}C(=O)NH_2$ ; wherein  $R^{2a}$ ,  $R^{2b}$  and  $R^{2c}$  is optionally substituted with at least one of halo, alkyl, alkylideneamine, arylideneamine, cyano, hydroxy, alkoxy, amino, amidino, guanidino, imino, amido, acid, ester, keto, aldehyde, dioxolane, furanyl, piperidinyl, piperazinyl, pyrrolidinyl, aryl, morpholinyl, and thiomorpholinyl dioxide; or  $R^{2a}$  and  $R^{2b}$  or  $R^{1a}$ , as set forth above, and  $R^{2a}$  or  $R^{1a}$ , as set forth above, and  $R^{2b}$  taken together with the nitrogen atom to which they are each attached can form a substituted or unsubstituted 3 to 8 membered heterocyclic or heteroaromatic amine group which, optionally, contains at least one other heteroatom of N, O or S; wherein  $R^{2a}$ ,  $R^{2b}$  or  $R^{2c}$  is optionally substituted with at least one of halo, alkyl, alkylideneamine, arylideneamine, cyano, hydroxy, alkoxy, amino, amidino, guanidino, imino, amido, acid, ester, keto, aldehyde, dioxolane, furanyl, piperidinyl, piperazinyl, pyrrolidinyl, aryl, morpholinyl, and thiomorpholinyl dioxide;


$R$  is, in each occurrence, independently, H,  $-C_{1-6}$  alkyl,  $-C_{3-8}$  cycloalkyl,  $-C_{0-6}$  alkyl- $OC_{1-6}$  alkyl,  $-C_{0-6}$  alkyl- $O(CH_2)_{1-4}-COOH$ ,  $-C_{0-6}$  alkyl- $O(CH_2)_{1-4}-C(=O)OC_{1-6}$  alkyl,  $-C_{0-6}$  alkyl $COOH$ ,  $-C_{0-6}$  alkyl $CO_2C_{1-6}$  alkyl,  $-C_{0-6}$  alkyl $OC_{1-6}$  alkyl,  $-C_{1-6}$  alkyl $OH$ ,  $-C_{0-6}$  alkyl $CONH_2$ ,  $-C_{0-6}$  alkyl $CONHC_{0-6}$  alkyl,  $-C_{0-6}$  alkyl $CON(C_{0-6}$  alky) $_2$ ,  $-C_{0-6}$  alkyl $CON(CH_2)_{2-6}$ ,  $-C_{0-6}$  alkyl $NH_2$ ,  $-C_{0-6}$  alkyl $NH(C_{1-6}$  alkyl) or  $-C_{0-6}$  alkyl $N(C_{1-6}$  alkyl) $_2$ .

25

Q is a member selected from the group consisting of:




16



$Y$  is S;

5  $R^1$  is H, -Cl, -Br, -I, -F, -OCF<sub>3</sub>, alkyl, hydroxy, alkoxy, amino, thiol, thioalkyl, thioaryl, or piperiziny;

$J^1$  is a member selected from the group consisting of:



10  $X$  is O or S;


$R^2$  is H, -Cl, -Br, -I, -F or -OC<sub>1-6</sub>alkyl;

$R^3$  is H, -Cl, -Br, -I, -F, -OC<sub>1-6</sub>alkyl, -NHC<sub>1-6</sub>acyl, -NO<sub>2</sub>, -NHSO<sub>2</sub>C<sub>1-4</sub>alkyl, -CN, -NH<sub>2</sub>, -CONH<sub>2</sub>, -SO<sub>2</sub>C<sub>1-6</sub>alkyl, -SO<sub>2</sub>NH<sub>2</sub>, -CO<sub>2</sub>C<sub>1-6</sub>alkyl or -O(CH<sub>2</sub>)<sub>1-4</sub>COOH;

15

$R^4$  and  $R^5$  are each independently H, -Cl, -Br, -I, -F or -OC<sub>1-6</sub>alkyl;

$J^2$  is a member selected from the group consisting of:



20

$Z$  is -NR<sup>6</sup>-, -O- or -S-;


R<sup>6</sup> is H, C<sub>1-6</sub>alkyl or C<sub>3-8</sub>cycloalkyl;

R<sup>7</sup> and R<sup>8</sup> are independently H, -Cl, -Br, -I or -F, where at least one of R<sup>7</sup> and R<sup>8</sup> is not  
5 hydrogen; and

R<sup>9</sup> and R<sup>10</sup> are independently H, -Cl, -Br, -I or -F, where at least one of R<sup>9</sup> and R<sup>10</sup> is not hydrogen;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and  
10 prodrug derivatives thereof.


The invention further provides a compound of the formulae (I) or (II):



wherein:

15


A is a member selected from the group consisting of:



R is, in each occurrence, independently, H, -C<sub>1-6</sub>alkyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>1-6</sub>alkylOH, -C<sub>0-6</sub>alkyl-OC<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkyl-O(CH<sub>2</sub>)<sub>1-4</sub>-COOH, -C<sub>0-6</sub>alkyl-O(CH<sub>2</sub>)<sub>1-4</sub>-C(=O)OC<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkylCOOH, -C<sub>0-6</sub>alkylCO<sub>2</sub>C<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkylCONH<sub>2</sub>,

-C<sub>0-6</sub>alkylCONHC<sub>0-6</sub>alkyl, -C<sub>0-6</sub>alkylCON(C<sub>0-6</sub>alkyl)<sub>2</sub>, -C<sub>0-6</sub>alkylCON(CH<sub>2</sub>)<sub>2-6</sub>, -C<sub>0-6</sub>alkylNH<sub>2</sub>, -C<sub>0-6</sub>alkylNH(C<sub>1-6</sub>alkyl) or -C<sub>0-6</sub>alkylN(C<sub>1-6</sub>alkyl)<sub>2</sub>.


Q is a member selected from the group consisting of:



5 Y is S;

R<sup>1</sup> is H, -Cl, -Br, -I or -F, -OMe, NH<sub>2</sub>, NHMe, NHMe<sub>2</sub>, -NHCOMe, -NHSO<sub>2</sub>Me;

J<sup>1</sup> is a member selected from the group consisting of:



X is O or S;

R<sup>2</sup> is H, -Cl, -Br, -I, -F or -OC<sub>1-6</sub>alkyl;


15

R<sup>3</sup> is H, -Cl, -Br, -I, -F, -OC<sub>1-6</sub>alkyl, -NHC<sub>1-6</sub>acyl, -NO<sub>2</sub>, -NHCO<sub>2</sub>C<sub>1-4</sub>alkyl, -CN or -O(CH<sub>2</sub>)<sub>1-4</sub>-COOH;

R<sup>4</sup> and R<sup>5</sup> are each independently H, -Cl, -Br, -I, -F or -OC<sub>1-6</sub>alkyl;

20

J<sup>2</sup> is a member selected from the group consisting of:



Z is  $\text{-NR}^6\text{-}$ ,  $\text{-O-}$  or  $\text{-S-}$ ;



5  $\text{R}^6$  is H,  $\text{C}_{1-6}$ alkyl or  $\text{C}_{3-8}$ cycloalkyl;

$\text{R}^7$  and  $\text{R}^8$  are each independently H, -Cl, -Br, -I or -F, where at least one of  $\text{R}^7$  and  $\text{R}^8$  is not hydrogen; and

10  $\text{R}^9$  and  $\text{R}^{10}$  are each independently H, -Cl, -Br, -I or -F, where at least one of  $\text{R}^9$  and  $\text{R}^{10}$  is not hydrogen;


and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

15 The invention further provides a compound of formula (I) or (II):




wherein:

20 A is a member selected from the group consisting of:



R is, in each occurrence, independently, H, -C<sub>1-6</sub>alkyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>1-6</sub>alkylOH, -C<sub>0-6</sub>alkyl-OC<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkyl-O(CH<sub>2</sub>)<sub>1-4</sub>-COOH, -C<sub>0-6</sub>alkyl-O(CH<sub>2</sub>)<sub>1-4</sub>-C(=O)OC<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkylCOOH, -C<sub>0-6</sub>alkylCO<sub>2</sub>C<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkylCONH<sub>2</sub>, -C<sub>0-6</sub>alkylCONHC<sub>0-6</sub>alkyl, -C<sub>0-6</sub>alkylCON(C<sub>0-6</sub>alkyl)<sub>2</sub>, -C<sub>0-6</sub>alkylCON(CH<sub>2</sub>)<sub>2-6</sub>, -C<sub>0-6</sub>alkylNH<sub>2</sub>, -C<sub>0-6</sub>alkylNH(C<sub>1-6</sub>alkyl) or -C<sub>0-6</sub>alkylN(C<sub>1-6</sub>alkyl)<sub>2</sub>.


Q is a member selected from the group consisting of:



Y is S;

R<sup>1</sup> is H, -Cl, -Br, -I or -F, -OMe, NH<sub>2</sub>, NHMe, NHMe<sub>2</sub>;

J<sup>1</sup> is a member selected from the group consisting of:




X is O or S;

R<sup>3</sup> is H, -Cl, -Br, -I or -F;

$R^5$  is H, -Cl, -Br, -I or -F;

$J^2$  is a member selected from the group consisting of:




$R^6$  is a H, C<sub>1-6</sub> alkyl or C<sub>3-8</sub> cycloalkyl;

10  $R^7$  and  $R^8$  are each independently -Cl, -Br, -I or -F; and

$R^9$  and  $R^{10}$  are each independently -Cl, -Br, -I or -F;


and all pharmaceutically acceptable isomers, salts, hydrates, solvates and  
15 prodrug derivatives thereof.


The invention provides compounds of formula (I) having the following structure:




wherein:

20 A is a member selected from the group consisting of:








5

10






and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug


15 derivatives thereof.












The invention further provides compounds of formula (I) having the following structure:



wherein:













5 A is a member selected from the group consisting of:






5

10




5

10




and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.


- 10 The invention further provides compounds of formula (I) having the following structure:



wherein:

Q is a member selected from the group consisting of:






5


and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug  
10 derivatives thereof.


The invention further provides compounds of formula (I) having the following  
structure:



wherein:

15 A is a member selected from the group consisting of:






5


10






5

10



and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug  
5 derivatives thereof.

The invention further provides compounds of formula (I) having the following structure:




wherein:

R is independently selected from the group consisting of:

H, -CO<sub>2</sub>H, -CO<sub>2</sub>Me, -CONH<sub>2</sub>, -CONHMe, -CONHMe<sub>2</sub>, -CON(CH<sub>2</sub>)<sub>4</sub>, -CON(CH<sub>2</sub>)<sub>5</sub>, -CH<sub>2</sub>OH, -CH<sub>2</sub>OMe, -CH<sub>2</sub>CO<sub>2</sub>H, -CH<sub>2</sub>CO<sub>2</sub>Me, -CH<sub>2</sub>CONH<sub>2</sub>, -CH<sub>2</sub>CH<sub>2</sub>OH, -CH<sub>2</sub>CH<sub>2</sub>OMe, -CH<sub>2</sub>NH<sub>2</sub>, -CH<sub>2</sub>N(Me)<sub>2</sub>, and -CH<sub>3</sub>,

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

The invention further provides compounds of formula (I) having the following structure:




wherein:

R is independently selected from the group consisting of:

H,  $-\text{CO}_2\text{H}$ ,  $-\text{CO}_2\text{Me}$ ,  $-\text{CONH}_2$ ,  $-\text{CONHMe}$ ,  $-\text{CONHMe}_2$ ,  $-\text{CON}(\text{CH}_2)_4$ ,  $-\text{CON}(\text{CH}_2)_5$ ,  $-\text{CH}_2\text{OH}$ ,  $-\text{CH}_2\text{OMe}$ ,  $-\text{CH}_2\text{CO}_2\text{H}$ ,  $-\text{CH}_2\text{CO}_2\text{Me}$ ,  $-\text{CH}_2\text{CONH}_2$ ,  $-\text{CH}_2\text{CH}_2\text{OH}$ ,  $-\text{CH}_2\text{CH}_2\text{OMe}$ ,  $-\text{CH}_2\text{NH}_2$ ,  $-\text{CH}_2\text{N}(\text{Me})_2$ , and  $-\text{CH}_3$ ,


and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug  
5 derivatives thereof.


The invention further provides compounds of formula (I) having the following  
structure:




wherein:

10 A is a member selected from the group consisting of:

















5




10



5


10



5

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug  
10 derivatives thereof.


The invention further provides compounds of formula (I) having the following structure:



15

wherein:


$J^2$  is a member selected from the group consisting of:



and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.


5

The invention further provides compounds of formula (II) having the following structure:



10 wherein:

J<sup>1</sup> is a member selected from the group consisting of:



and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

- 5 The invention also encompasses all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives of the compounds of the invention as set forth herein. The compounds of the invention can exist in various isomeric and tautomeric forms, and all such forms are meant to be included in the invention, along with pharmaceutically acceptable salts, hydrates, solvates and prodrug derivatives of
- 10 such isomers and tautomers.

The compounds of the invention may be isolated as the free acid or base or converted to salts of various inorganic and organic acids and bases. Such salts are



within the scope of the invention. Non-toxic and physiologically compatible salts are particularly useful although other less desirable salts may have use in the processes of isolation and purification.

A number of methods are useful for the preparation of the salts of the  
5 compounds as described above and are known to those skilled in the art. For example, the free acid or free base form of a compound of one of the formulae above can be reacted with one or more molar equivalents of the desired acid or base in a solvent or solvent mixture in which the salt is insoluble, or in a solvent like water after which the solvent is removed by evaporation, distillation or freeze drying. Alternatively, the free  
10 acid or base form of the product may be passed over an ion exchange resin to form the desired salt or one salt form of the product may be converted to another using the same general process.

The invention also encompasses prodrug derivatives of the compounds contained herein. The term "prodrug" refers to a pharmacologically inactive derivative  
15 of a parent drug molecule that requires biotransformation, either spontaneous or enzymatic, within the organism to release the active drug. Prodrugs are variations or derivatives of the compounds of the invention which have groups cleavable under metabolic conditions. Prodrugs become the compounds of the invention which are pharmaceutically active *in vivo*, when they undergo solvolysis under physiological  
20 conditions or undergo enzymatic degradation. Prodrug compounds of the invention may be called single, double, triple etc., depending on the number of biotransformation steps required to release the active drug within the organism, and indicating the number of functionalities present in a precursor-type form. Prodrug forms often offer advantages of solubility, tissue compatibility, or delayed release in the mammalian  
25 organism (see, Bundgard, Design of Prodrugs, pp. 7-9, 21-24, Elsevier, Amsterdam 1985 and Silverman, The Organic Chemistry of Drug Design and Drug Action, pp. 352-401, Academic Press, San Diego, CA, 1992). Prodrugs commonly known in the art include acid derivatives well known to practitioners of the art, such as, for example, esters prepared by reaction of the parent acids with a suitable alcohol, or amides  
30 prepared by reaction of the parent acid compound with an amine, or basic groups

reacted to form an acylated base derivative. Moreover, the prodrug derivatives of the invention may be combined with other features herein taught to enhance bioavailability.

The compounds of the present invention may also be used alone or in combination or in combination with other therapeutic or diagnostic agents. In certain preferred embodiments, the compounds of the invention may be coadministered along with other compounds typically prescribed for these conditions according to generally accepted medical practice such as anticoagulant agents, thrombolytic agents, or other antithrombotics, including platelet aggregation inhibitors, tissue plasminogen activators, urokinase, prourokinase, streptokinase, heparin, aspirin, or warfarin. The compounds of the present invention may act in a synergistic fashion to prevent reocclusion following a successful thrombolytic therapy and/or reduce the time to reperfusion. These compounds may also allow for reduced doses of the thrombolytic agents to be used and therefore minimize potential hemorrhagic side-effects. The compounds of the invention can be utilized *in vivo*, ordinarily in mammals such as primates (e.g. humans), sheep, horses, cattle, pigs, dogs, cats, rats and mice, or *in vitro*.

The biological properties of the compounds of the present invention can be readily characterized by methods that are well known in the art, for example by the *in vitro* protease activity assays and *in vivo* studies to evaluate antithrombotic efficacy, and effects on hemostasis and hematological parameters, such as are illustrated in the examples.

Diagnostic applications of the compounds of the invention will typically utilize formulations in the form of solutions or suspensions. In the management of thrombotic disorders the compounds of the invention may be utilized in compositions such as tablets, capsules or elixirs for oral administration, suppositories, sterile solutions or suspensions or injectable administration, and the like, or incorporated into shaped articles. Subjects in need of treatment (typically mammalian) using the compounds of the invention can be administered dosages that will provide optimal efficacy. The dose and method of administration will vary from subject to subject and be dependent upon such factors as the type of mammal being treated, its sex, weight, diet, concurrent medication, overall clinical condition, the particular compounds employed, the specific use for which these compounds are employed, and other factors which those skilled in

the medical arts will recognize.

#### Preparation of Compounds

The compounds of the present invention may be synthesized by standard  
5 organic chemical synthetic methods as described and referenced in standard textbooks. These methods are well known in the art. See, e.g., Morrison and Boyd, "Organic Chemistry", Allyn and Bacon, Inc., Boston, 1959, et seq.


Starting materials used in any of these methods are commercially available from  
chemical vendors such as Aldrich, Sigma, Nova Biochemicals, Bachem Biosciences,  
10 and the like, or may be readily synthesized by known procedures.

Reactions are carried out in standard laboratory glassware and reaction vessels  
under reaction conditions of standard temperature and pressure, except where otherwise  
indicated.


During the synthesis of these compounds, the functional groups of the  
15 substituents are optionally protected by blocking groups to prevent cross reaction during coupling procedures. Examples of suitable blocking groups and their use are described in "The Peptides: Analysis, Synthesis, Biology", Academic Press, Vol. 3 (Gross, *et al.*, Eds., 1981) and Vol. 9 (1987), the disclosures of which are incorporated herein by reference.

20 Non-limiting exemplary synthesis schemes are outlined directly below, and specific steps are described in the Examples. The reaction products are isolated and purified by conventional methods, typically by solvent extraction into a compatible solvent. The products may be further purified by column chromatography or other appropriate methods.

## SCHEME 1




## SCHEME 2




5

## SCHEME 3




10

## SCHEME 4




## SCHEME 5




5

## SCHEME 6




## SCHEME 7




10


## SCHEME 8




## SCHEME 9



## SCHEME 10




## SCHEME 11



The following examples are non-limiting embodiments of the present invention,  
 5 which were made utilizing a method as generally shown in reaction Schemes 1-11  
 above or by a similar procedure as would be understood by one of skill in the art.

ExamplesExample 1

- 10 **Ethyl 2-[4-({4-[(dimethylamino)iminomethyl]phenyl}methyl)-1-[2-(5-chloro(2-thienyloxy))acetyl]-3-oxopiperazin-2-yl]acetate.**



Part 1. To a solution of ethyl 3-oxopiperazine-2-acetate (482.7 mg, 2.59 mmol) in DMF (8 mL) at room temperature was added 5-chloro-2-thienyloxyacetic acid (Ewing, W. R., WO 0032590, 2000) (415 mg, 2.16 mmol), DIEA (0.75 mL, 4.32 mmol), and BOP (1.15 g, 2.59 mmol). The solution was stirred at room temperature for 24 hours. The reaction mixture was diluted in EtOAc and water. The organic layer was washed with sat. NaHCO<sub>3</sub> and sat. NaCl, dried over MgSO<sub>4</sub>, filtered and concentrated *in vacuo* to yield ethyl 2-{1-[2-(5-chloro(2-thienyloxy))acetyl]-3-oxopiperazin-2-yl}acetate (0.56 g, 72% yield). MS found for (M+H)<sup>+</sup>: 361.1.


Part 2. To a solution ethyl 2-{1-[2-(5-chloro(2-thienyloxy))acetyl]-3-oxopiperazin-2-yl}acetate (560 mg, 1.56 mmol) and -bromo-p-tolunitrile (305 mg, 1.56 mmol) in 5 mL of DMF was added Cs<sub>2</sub>CO<sub>3</sub> (762 mg, 2.34 mmol). After stirring at rt for 24 h, the mixture was diluted with EtOAc and washed with H<sub>2</sub>O. The organic layer was dried over MgSO<sub>4</sub>, filtered and evaporated *in vacuo*. Flash chromatography on silica gel (10% EtOAc in DCM) gave ethyl 2-{1-[2-(5-chloro(2-thienyloxy))acetyl]-4-[(4-cyanophenyl)methyl]-3-oxopiperazin-2-yl}acetate (320 mg, 43%). MS found for C<sub>13</sub>H<sub>9</sub>N<sub>2</sub>O<sub>3</sub> (M+H)<sup>+</sup>: 476.

Part 3. A stream of H<sub>2</sub>S (g) was bubbled through a solution of ethyl 2-{1-[2-(5-chloro(2-thienyloxy))acetyl]-4-[(4-cyanophenyl)methyl]-3-oxopiperazin-2-yl}acetate (320 mg, 0.67 mmol) in 4.5 mL pyridine and 0.5 mL NEt<sub>3</sub> until saturation. The mixture was stirred at rt for 4-5 hr and evaporated. The resulting residue was treated with MeI (0.42 mL, 6.74 mmol) in 5 mL acetone at reflux temperature for 1 hr and concentrated to dryness. The resulting residue was treated with a mixture of dimethylamine (1.68 mL of 2 M solution in THF, 3.37 mmol) in acetic acid (0.29 mL, 5.1 mmol) and 4 mL methanol under reflux for 30min. The solvent was removed at reduced pressure and the crude product was purified by HPLC (C18 reversed phase) eluting with 0.1% TFA in H<sub>2</sub>O/CH<sub>3</sub>CN to give Ethyl 2-[4-({4-[(dimethylamino)iminomethyl]phenyl}methyl)-1-[2-(5-chloro(2-thienyloxy))acetyl]-3-oxopiperazin-2-yl]acetate. MS found: (M+H)<sup>+</sup>: 521.1.



Example 2


**Ethyl 2-(1-[2-(5-chloro(2-thienyloxy))acetyl]-4-{{4-(iminopyrrolidinylmethyl)phenyl}methyl}-3-oxopiperazin-2-yl)acetate.**



- 5 The titled compound was synthesized using a similar procedure to that described in Example 1 using pyrrolidine instead of dimethylamine in Part 3. MS found: (M+H)<sup>+</sup>: 580.1.

Example 3


- 10 **2-[4-({4-[(Dimethylamino)iminomethyl]phenyl}methyl)-1-[2-(5-chloro(2-thienyloxy))acetyl]-3-oxopiperazin-2-yl]acetic acid.**



- A solution of ethyl 2-[4-({4-[(dimethylamino)iminomethyl]phenyl}methyl)-1-[2-(5-chloro(2-thienyloxy))acetyl]-3-oxopiperazin-2-yl]acetate (40 mg, 0.077 mmol) in 2 mL of methanol and 2 mL of H<sub>2</sub>O was treated with LiOH (6.46 mg, 0.154 mmol) at rt for 2 h. Methanol was evaporated, and the H<sub>2</sub>O layer was acidified with 1N HCl until pH ~ 1-2. The H<sub>2</sub>O layer was extracted with EtOAc. The organic layer was dried over MgSO<sub>4</sub>, filtered and evaporated *in vacuo* and the crude product was purified by HPLC (C18 reversed phase) eluting with 0.1% TFA in H<sub>2</sub>O/CH<sub>3</sub>CN to give 2-[4-({4-[(Dimethylamino)iminomethyl]phenyl}methyl)-1-[2-(5-chloro(2-thienyloxy))acetyl]-3-oxopiperazin-2-yl]acetic acid. MS found: (M+H)<sup>+</sup>: 493.1.

Example 4


- 25 **2-(1-[2-(5-Chloro(2-thienyloxy))acetyl]-4-{{4-(iminopyrrolidinylmethyl)phenyl}methyl}-3-oxopiperazin-2-yl)acetic acid.**



The titled compound was synthesized using a similar procedure to that described in Example 3 using ethyl 2-(1-[2-(5-chloro(2-thienyloxy))acetyl]-4-[[4-(iminopyrrolidinylmethyl)phenyl]methyl]-3-oxopiperazin-2-yl)acetate. MS found: 5 (M+H)<sup>+</sup>: 579.1.

#### Example 5

**Ethyl 2-[4-({4-[(dimethylamino)iminomethyl]phenyl}methyl)-1-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-3-oxopiperazin-2-yl]acetate.**



10


Part 1. To a solution of ethyl 3-oxopiperazine-2-acetate (1.12 g, 6 mmol) in pyridine (15 mL) at rt, was added 6-chlorobenzo[b]thiophene-2-sulfonyl chloride (Ewing, W. R., WO 9937304) (1.33 g, 5 mmol). The mixture was stirred at rt overnight and evaporated *in vacuo*. The residue was dissolved in EtOAc and H<sub>2</sub>O. The organic layer 15 was washed with water, 1N HCl, sat. NaHCO<sub>3</sub>, sat. NaCl, dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated to give ethyl 2-{1-[(6-chlorobenzo[b]thiophen-2-yloxy)sulfinyl]-3-oxopiperazin-2-yl}acetate as a solid (1.43 g, 69% yield). MS found (M+H)<sup>+</sup>: 417.1.

Part 2. To a solution of ethyl 2-{1-[(6-chlorobenzo[b]thiophen-2-yloxy)sulfinyl]-3-oxopiperazin-2-yl}acetate (1.04 g, 2.5 mmol) and -bromo-p-tolunitrile (588mg, 3 mmol) in 15 mL of DMF was added Cs<sub>2</sub>CO<sub>3</sub> (1.22 g, 3.75 mmol). After stirring at rt for 24 h, the mixture was diluted with EtOAc and washed with H<sub>2</sub>O. The organic layer was dried over MgSO<sub>4</sub>, filtered and evaporated *in vacuo*. Flash chromatography on silica gel (40% EtOAc in hexane) gave ethyl 2-{1-[(6-chlorobenzo[b]thiophen-2-yloxy)sulfinyl]-4-[(4-cyanophenyl)methyl]-3-oxopiperazin-2-yl}acetate (320 mg, 25 43%). MS found (M+H)<sup>+</sup>: 532.1.

Part 3. A stream of H<sub>2</sub>S (g) was bubbled through a solution of ethyl 2-{1-[(6-chlorobenzo[b]thiophen-2-yloxy)sulfinyl]-4-[(4-cyanophenyl)methyl]-3-oxopiperazin-2-yl}acetate (524 mg, 0.985 mmol) in 9 mL pyridine and 1 mL NEt<sub>3</sub> until saturation. The mixture was stirred at rt for 24 hr and evaporated. The resulting  
 5 residue was treated with MeI (0.613 mL, 9.85 mmol) in 5 mL acetone at reflux temperature for 1 hr and concentrated to dryness. The resulting residue was treated with a mixture of dimethylamine (2.48 mL of 2 M solution in THF, 4.92 mmol) in acetic acid (0.42 mL, 7.38 mmol) and 16 mL methanol under reflux for 30min. The solvent was removed at reduced pressure and the crude product was purified by HPLC  
 10 (C18 reversed phase) eluting with 0.1% TFA in H<sub>2</sub>O/CH<sub>3</sub>CN to give ethyl 2-[4-({4-[(dimethylamino)iminomethyl]phenyl}methyl)-1-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-3-oxopiperazin-2-yl]acetate. MS found: (M+H)<sup>+</sup>: 577.1.

#### Example 6

15 Ethyl 2-(1-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-4-{[4-(iminopyrrolidinylmethyl)phenyl]methyl}-3-oxopiperazin-2-yl)acetate.




The titled compound was synthesized using a similar procedure to that described in Example 5 using pyrrolidine instead of dimethylamine in Part 3. MS found: (M+H)<sup>+</sup>:  
 20 604.1.

#### Example 7

Ethyl 2-(1-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-4-{[4-(1-methyl(2-imidazol-2-yl))phenyl]methyl}-3-oxopiperazin-2-yl)acetate.


25



The titled compound was synthesized using a similar procedure to that described in Example 5 using N-methylethylenediamine instead of dimethylamine in Part 3. MS found: (M+H)<sup>+</sup>: 590.1.

5 **Example 8**


**2-[4-({4-[(dimethylamino)iminomethyl]phenyl}methyl)-1-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-3-oxopiperazin-2-yl]acetic acid.**



- 10 A solution of ethyl 2-[4-({4-[(dimethylamino)iminomethyl]phenyl}methyl)-1-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-3-oxopiperazin-2-yl]acetate (40 mg, 0.07 mmol) in 2 mL of methanol and 2 mL of H<sub>2</sub>O was treated with LiOH (5.8 mg, 0.14 mmol) at rt for 2 h. Methanol was evaporated, and the H<sub>2</sub>O layer was acidified with 1N HCl until pH ~ 1-2. The H<sub>2</sub>O layer was extracted with EtOAc. The organic layer was dried over
- 15 MgSO<sub>4</sub>, filtered and evaporated *in vacuo* and the crude product was purified by HPLC (C18 reversed phase) eluting with 0.1% TFA in H<sub>2</sub>O/CH<sub>3</sub>CN to give 2-[4-({4-[(dimethylamino)iminomethyl]phenyl}methyl)-1-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-3-oxopiperazin-2-yl]acetic acid. Mass found: (M+H)<sup>+</sup>: 550.

20 **Example 9**


**2-(1-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-4-{[4-(iminopyrrolidinylmethyl)phenyl]methyl}-3-oxopiperazin-2-yl)acetic acid.**



- The titled compound was synthesized using a similar procedure to that described in
- 25 Example 8 using Ethyl 2-(1-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-4-{[4-(iminopyrrolidinylmethyl)phenyl]methyl}-3-oxopiperazin-2-yl)acetate. MS found: (M+H)<sup>+</sup>: 576.1.

Example 10

2-(1-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-4-{[4-(1-methyl(2-imidazolin-2-yl))phenyl]methyl}-3-oxopiperazin-2-yl)acetic acid.



5


The titled compound was synthesized using a similar procedure to that described in Example 8 using ethyl 2-(1-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-4-{[4-(1-methyl(2-imidazolin-2-yl))phenyl]methyl}-3-oxopiperazin-2-yl)acetate.

MS found: (M+H)<sup>+</sup>: 562.1.

10

Example 11

1-({4-[(Dimethylamino)iminomethyl]phenyl}methyl)-4-[2-(4-chlorophenoxy)acetyl]piperazin-2-one.



- 15 Part 1. A mixture of 4-benzyloxycarbonylpiperazin-2-one (3 g, 12.82 mmol), -bromo-p-tolunitrile (6.71 g, 19.23 mmol), and Cs<sub>2</sub>CO<sub>3</sub> (12.53 g, 38.46 mmol) in DMF (10 mL) was stirred at 50°C for 24 hours. The solid was filtered and the filtrate was concentrated. The residue was diluted with EtOAc, washed with sat. NaCl (2x), dried and evaporated. The crude material was subjected to silica gel chromatography to
- 20 afford phenylmethyl 4-[(4-cyanophenyl)methyl]-3-oxopiperazinecarboxylate (3.5 g, 78%). MS found: (M+H)<sup>+</sup>: 350.1.

Part 2. A mixture of phenylmethyl 4-[(4-cyanophenyl)methyl]-3-

oxopiperazinecarboxylate (1 g, 2.87 mmol) and 10% Pd/C (160 mg) in MeOH (10 mL)

- 25 was stirred under balloon H<sub>2</sub> (1 atm) for 3 hrs. The Pd/C was filtered off through a celite bed. The filtrate was evaporated to give 4-[(2-oxopiperazinyl)methyl]benzenecarbonitrile (600 mg, 97%). MS found: (M+H)<sup>+</sup>: 216.1.

Part 3. A mixture of 4-chlorophenol (3 g, 23.44 mmol), tert-butyl bromoacetate (6.82 g, 35.16 mmol), and  $\text{Cs}_2\text{CO}_3$  (23.04 g, 70.72 mmol) in DMF (50 mL) was stirred at 50°C for 24 hours. The solid was filtered and the filtrate was concentrated. The residue was  
5 diluted with EtOAc and water, washed with sat. NaCl (2x), dried and evaporated. The crude material was subjected to silica gel chromatography to afford tert-butyl 2-(4-chlorophenoxy)acetate (5.2 g, 92%). MS found: (M+H)<sup>+</sup>: 243.1.

Part 4. tert-butyl 2-(4-chlorophenoxy)acetate (2 g, 8.26 mmol) was dissolved in  
10 dioxane. 6 M HCl (10 mL) was added. The mixture was stirred at rt for 1 hr and diluted with EtOAc (20 mL) and water (10 mL). The organic layer was separated and extracted into 0.5 N NaOH (10 mL). H<sub>2</sub>O layer was washed with EtOAc and acidified with 1N HCl until pH ~ 1-2. The H<sub>2</sub>O layer was extracted with EtOAc. The organic layer was dried over  $\text{MgSO}_4$ , filtered and evaporated *in vacuo*. Thionyl chloride (5 mL) was  
15 added to the residue and the mixture was refluxed for 2 hrs, evaporated in *vacuo* to give 2-(4-chlorophenoxy)acetyl chloride (1.34 g, 80% yield). MS found: (M+H)<sup>+</sup>: 205.1.


Part 5. To a solution of 4-[(2-oxopiperaziny)methyl]benzenecarbonitrile (600 mg, 2.79 mmol) and 2-(4-chlorophenoxy)acetyl chloride (683 mg, 3.35 mmol) in  $\text{CH}_3\text{CN}$  (10  
20 mL) at rt, was added N-methylmorpholine (1.13 g, 11.16 mmol). The mixture was stirred at rt for 12 hrs. The solvent was evaporated. The residue was diluted with EtOAc, washed with 1 N HCl, sat.  $\text{NaHCO}_3$ , sat. NaCl, dried with  $\text{Na}_2\text{SO}_4$  and evaporated to afford 4-({4-[2-(4-chlorophenoxy)acetyl]-2-oxopiperaziny}methyl)benzenecarbonitrile (750 mg, 70% yield). MS found: (M+H)<sup>+</sup>:  
25 394.1.

Part 6. A stream of HCl (g) was bubbled through a solution of part 5 (400 mg, 1.04 mmol) in MeOH (10 mL) at 0°C for 10 min. The mixture was stirred at rt for 4 hr and evaporated in *vacuo*. The resulting residue was dissolved in MeOH (30 mL), treated  
30 with dimethylamine (2.61 mL of 2 M solution in THF, 5.22 mmol) at reflux temperature for 1 hr and concentrated to dryness. The crude product was purified by

HPLC (C18 reversed phase) eluting with 0.1% TFA in H<sub>2</sub>O/CH<sub>3</sub>CN to give -({4-[(Dimethylamino)iminomethyl]phenyl}methyl)-4-[2-(4-chlorophenoxy)acetyl]piperazin-2-one. MS found: (M+H)<sup>+</sup>: 429.1.

### 5 Example 12


**4-[2-(4-Chlorophenoxy)acetyl]-1-{[4-(1-methyl(2-imidazolin-2-yl))phenyl]methyl}piperazin-2-one.**



The titled compound was synthesized using a similar procedure to that described in  
 10 Example 11 using N-methylethylenediamine instead of dimethylamine in part 6. MS found: (M+H)<sup>+</sup>: 441.1.

### Example 13

**1-{[4-(Azetidinyliminomethyl)phenyl]methyl}-4-[2-(4-chlorophenoxy)acetyl]piperazin-2-one.**




The titled compound was synthesized using a similar procedure to that described in  
 Example 11 using azetidine instead of dimethylamine in Part 6. MS found:  
 (M+H)<sup>+</sup>:441.1.

20

### Example 14


**4-[2-(4-Chlorophenoxy)acetyl]-1-{[4-(iminopyrrolidinylmethyl)phenyl]methyl}piperazin-2-one.**



The titled compound was synthesized using a similar procedure to that described in Example 11 using pyrrolidine instead of dimethylamine in Part 6. MS found: (M+H)<sup>+</sup>: 455.1.

### 5 Example 15


**4-[2-(4-Chlorophenoxy)acetyl]-1-[[4-(iminopiperidylmethyl)phenyl]methyl]piperazin-2-one.**



The titled compound was synthesized using a similar procedure to that described in Example 11 using piperidine instead of dimethylamine in Part 6. MS found: (M+H)<sup>+</sup>: 469.1.

### Example 16

**1-({4-[(Dimethylamino)iminomethyl]phenyl}methyl)-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]piperazin-2-one.**



Part 1. To a solution of phenylmethyl 4-[(4-cyanophenyl)methyl]-3-oxopiperazinecarboxylate (1.29 g, 6 mmol) in pyridine (15 mL) at rt, was added 6-chlorobenzo[b]thiophene-2-sulfonyl chloride (1.33 g, 5 mmol). The mixture was stirred at rt overnight and evaporated *in vacuo*. The residue was dissolved in EtOAc and H<sub>2</sub>O. The organic layer was washed with water, 1N HCl, sat. NaHCO<sub>3</sub>, sat. NaCl, dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated to give 4-({4-[(6-chlorobenzo[b]thiophen-2-yloxy)sulfinyl]-2-oxopiperazinyl}methyl)benzenecarbonitrile as a solid (1.54 g, 69% yield). MS found (M+H)<sup>+</sup>: 446.1.

25


Part 2. A stream of H<sub>2</sub>S (g) was bubbled through a solution of 4-({4-[(6-chlorobenzo[b]thiophen-2-yloxy)sulfinyl]-2-



oxopiperazinyl)methyl)benzenecarbonitrile (438 mg, 0.985 mmol) in 9 mL pyridine and 1 mL  $\text{NEt}_3$  until saturation. The mixture was stirred at rt for 24 hr and evaporated. The resulting residue was treated with MeI (0.613 mL, 9.85 mmol) in 5 mL acetone at reflux temperature for 1 hr and concentrated to dryness. The resulting residue was  
 5 treated with a mixture of dimethylamine (2.48 mL of 2 M solution in THF, 4.92 mmol) in acetic acid (0.42 mL, 7.38 mmol) and 16 mL methanol at rt for 30 min. The solvent was removed at reduced pressure and the crude product was purified by HPLC (C18 reversed phase) eluting with 0.1% TFA in  $\text{H}_2\text{O}/\text{CH}_3\text{CN}$  to give 1-({4-  
 [(Dimethylamino)iminomethyl]phenyl)methyl}-4-[(6-chlorobenzo[b]thiophen-2-  
 10 yl)sulfonyl]piperazin-2-one. MS found:  $(\text{M}+\text{H})^+$ :491.1.

#### Example 17

**4-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-{[4-(1-methyl(2-imidazolin-2-yl))phenyl]methyl}piperazin-2-one.**




15

The titled compound was synthesized using a similar procedure to that described in Example 16 using N-methylethylenediamine instead of dimethylamine in Part 2. MS found:  $(\text{M}+\text{H})^+$ : 503.1.

#### 20 Example 18


**1-{{4-(Azetidinyloiminomethyl)phenyl)methyl}-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]piperazin-2-one.**



The titled compound was synthesized using a similar procedure to that described in  
 25 Example 16 using azetidine instead of dimethylamine in Part 2. MS found:  $(\text{M}+\text{H})^+$ : 503.1.

#### Example 19


4-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-[[4-(iminopyrrolidinylmethyl)phenyl]methyl]piperazin-2-one.



The titled compound was synthesized using a similar procedure to that described in Example 16 using pyrrolidine instead of dimethylamine in Part 2. MS found: (M+H)<sup>+</sup>: 517.1.

#### Example 20

Ethyl 1-[[4-({4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-2-oxopiperazinyl}methyl)phenyl]iminomethyl]piperidine-4-carboxylate.




The titled compound was synthesized using a similar procedure to that described in Example 16 using ethyl isonipecotat instead of dimethylamine in Part 2. MS found: (M+H)<sup>+</sup>: 603.1.

15

#### Example 21

1-[[4-({4-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-2-oxopiperazinyl}methyl)phenyl]iminomethyl]piperidine-4-carboxylic acid.



20 A solution of Ethyl 1-[[4-({4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-2-oxopiperazinyl}methyl)phenyl]iminomethyl]piperidine-4-carboxylate.

(46 mg, 0.077 mmol) in 2 mL of methanol and 2 mL of H<sub>2</sub>O was treated with LiOH (6.46 mg, 0.154 mmol) at rt for 2 h. Methanol was evaporated, and the H<sub>2</sub>O layer was acidified with 1N HCl until pH ~ 1-2. The H<sub>2</sub>O layer was extracted with EtOAc. The organic layer was dried over MgSO<sub>4</sub>, filtered and evaporated *in vacuo* and the crude


25

product was purified by HPLC (C18 reversed phase) eluting with 0.1% TFA in H<sub>2</sub>O/CH<sub>3</sub>CN to give 1-{{4-({4-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-2-oxopiperazinyl)methyl}phenyl)iminomethyl}piperidine-4-carboxylic acid.  
MS found: (M+H)<sup>+</sup>: 575.1.

5

#### Example 22

**1-{{4-(Azetidinyliminomethyl)phenyl)methyl}-4-[(5-chloroindol-2-yl)sulfonyl]piperazin-2-one.**



- 10 Part 1: To a 25 mL round bottom flask was added phenylmethyl 3-oxopiperazinecarboxylate (1.17 g, 5 mmol), dry DMF (10 mL), -bromo-p-tolunitrile (1 g, 5 mmol) and Cs<sub>2</sub>CO<sub>3</sub> (4.89 g, 15 mmol). The mixture was stirred at room temperature for 20 hours. EtOAc and water were added to the reaction. The organic layer was washed with water, dried over MgSO<sub>4</sub>, and concentrated *in vacuo* to afford  
15 phenylmethyl 4-[(4-cyanophenyl)methyl]-3-oxopiperazinecarboxylate as light yellow organic oil (1.43 g, 82%). MS found: (M+H)<sup>+</sup>: 350.

Part 2. A mixture of phenylmethyl 4-[(4-cyanophenyl)methyl]-3-oxopiperazinecarboxylate (1 g, 2.87 mmol) and 10% Pd/C (160 mg) in MeOH (10 mL)  
20 was stirred under balloon H<sub>2</sub> (1 atm) for 3 hrs. The Pd/C was filtered off through a celite bed. The filtrate was evaporated to give 4-[(2-oxopiperazinyl)methyl]benzenecarbonitrile (600 mg, 97%). MS found: (M+H)<sup>+</sup>: 216.1.

Part 3: A solution of 5-chloroindole (25.7 g, 169 mmol) in anhydrous THF (500 mL)  
25 was cooled with a dry ice-acetone bath, and n-BuLi (80 mL of 2.45 M solution in hexanes) was added dropwise over 15 minutes. The reaction was stirred in the cold for 10 minutes, then a solution of di-*t*-butyl dicarbonate (46.2 g, 212 mmol) in THF (150 mL) was added dropwise over 20 minutes. The reaction was stirred at room temperature overnight. EtOAc (500 mL) and 50 mL of water (50 mL) were added. The

aqueous layer was extracted with EtOAc (300 mL), and the combined organics were washed with water (200 mL), dried over MgSO<sub>4</sub>, filtered and the Filtrate were concentrated to give brown oil. This oil was subjected to flash column chromatography on silica gel first with 100% hexane, then with 0.5% EtOAc/Hexanes then with 1%  
5 EtOAc/Hexanes as eluents. The appropriate fractions were combined and concentrated to give tert-butyl 5-chloroindolecarboxylate (39.34 g, 92%) as clear, nearly colorless oil that crystallized on the high vacuum pump overnight.

Part 4: A solution of tert-butyl 5-chloroindolecarboxylate (20.0 g, 79.5 mmol) from  
10 Part 1 in THF (200 mL) was cooled with a dry ice-acetone bath under Ar, and n-BuLi (44 mL of 2.0 M solution in hexanes, 87.5 mmol) was added over 15 minutes. The reaction mixture was allowed to stir in the cold for 10 minutes, then added dropwise (over 10 minutes) via double-ended needle to a pre-cooled (with a dry ice-acetone bath) solution of SO<sub>2</sub> (80 g) in THF (100 mL). The reaction was allowed to stir at room  
15 temperature for 2 hrs. then concentrated to give a brown foam. This foam was dissolved in methylene chloride (200 mL) and the solution was cooled to 0C in an ice-water bath. To this was added sulfuryl chloride (7 mL, 87.5 mmol) dropwise over 5 minutes. The ice bath was removed and the reaction was stirred at room temperature overnight. Much solid (LiCl) had begun to form during the addition of SO<sub>2</sub>Cl<sub>2</sub>. The  
20 reaction was filtered and the filtrate was concentrated to give a dark residue. This residue was washed through a plug of silica gel with 1L of 30% DCM/Hexanes. The solvent was evaporated and the residue was subjected to flash column chromatography on silica gel using first 10% DCM/Hexanes, then 15% DCM/Hexanes. The appropriate fractions were combined and concentrated to give tert-butyl 5-chloro-2-  
25 (chlorosulfonyl)indolecarboxylate (14.34 g, 52%) as an off-white solid.

Part 5: Phenylmethyl 4-[(4-cyanophenyl)methyl]-3-oxopiperazinecarboxylate from Part 2 (0.35 g, 1 mmol) was dissolved in methylene chloride (5 mL) and pyridine (5 mL). Five minutes later 0.3502 g of tert-butyl 5-chloro-2-(chlorosulfonyl)indolecarboxylate  
30 (0.35 g) from Part 4 was added to the reaction. The mixture was stirred at room temperature for five hours. The solvent was removed *in vacuo*. Water and methylene

chloride was added to the crude brown oil. The organic layers were combined, dried over  $\text{MgSO}_4$ , concentrated *in vacuo* to *tert*-butyl 5-chloro-2-({4-[(4-cyanophenyl)methyl]-3-oxopiperazinyl}sulfinyloxy)indolecarboxylate as a light yellow solid (0.40 g, 75%). MS found:  $(\text{M}+\text{H})^+$ : 530.


5

Part 6: *tert*-Butyl 5-chloro-2-({4-[(4-cyanophenyl)methyl]-3-oxopiperazinyl}sulfinyloxy)indolecarboxylate from Part 5 (50 mg, 0.09 mmol) was dissolved in dry EtOH (10 mL) and HCl gas was bubbled through the solution until saturation. The mixture was stirred at room temperature for 24 hours. The solvent was removed *in vacuo* and the residue was redissolved in EtOH, to this was added azetidine HCl salt (50 mg, 0.53 mmol) and triethyl amine (84 mM, 0.6 mmol). The mixture was stirred at room temperature for 5 hours and purified by HPLC (C18 reversed phase) eluting with 0.1% TFA in  $\text{H}_2\text{O}/\text{CH}_3\text{CN}$  to afford 1-{{4-(Azetidyniliminomethyl)phenyl)methyl}-4-[(5-chloroindol-2-yl)sulfonyl]piperazin-2-one (34 mg, 74%). MS found  $(\text{M}+\text{H})^+$ : 486.

15

#### Example 23

Ethyl 1-{{4-[(4-[(5-chloroindol-2-yl)sulfonyl]-2-oxopiperazinyl)methyl]phenyl}iminomethyl}piperidine-4-carboxylate.




20

The titled compound was synthesized using a similar procedure to that described in Example 22 using ethyl isonipecotate instead of azetidine in Part 6. MS found:  $(\text{M}+\text{H})^+$ : 586.

#### 25 Example 24

1-({4-[(8-Aza-1,4-dioxaspiro[4.5]dec-8-yl)iminomethyl]phenyl)methyl}-4-[(5-chloroindol-2-yl)sulfonyl]piperazin-2-one.




The titled compound was synthesized using a similar procedure to that described in Example 22 using 1, 4-dioxo-8-azaspiro[4, 4]decane instead of azetidine in Part 6. MS found:  $(M+H)^+$ : 572.

5

#### Example 25


**4-[(5-chloroindol-2-yl)sulfonyl]-1-({4-[(ethylmethylamino)iminomethyl]phenyl}methyl)piperazin-2-one.**



10 The titled compound was synthesized using a similar procedure to that described in Example 22 using N-ethylmethylamine instead of azetidine in Part 6. MS found:  $(M+H)^+$ : 488.

#### Example 26


15 **1-[(4-{4-(Dimethylamino)piperidyl}iminomethyl}phenyl)methyl]-4-[(6-chloroindol-2-yl)sulfonyl]piperazin-2-one.**



The titled compound was synthesized using a similar procedure to that described in Example 22 using 4-(dimethylamino)-piperidine instead of azetidine in Part 6. MS  
20 found:  $(M+H)^+$ : 557.

#### Example 27

**1-({4-[(Dimethylamino)iminomethyl]phenyl}methyl)-4-[(6-chloroindol-2-yl)sulfonyl]piperazin-2-one.**




The titled compound was synthesized using a similar procedure to that described in Example 22 using dimethylamine instead of azetidine in part 6. MS found: (M+H)<sup>+</sup>: 474.

5

#### Example 28


**4-[(6-Chloroindol-2-yl)sulfonyl]-1-[[4-(iminopiperidylmethyl)phenyl]methyl]piperazin-2-one.**



10 The titled compound was synthesized using a similar procedure to that described in Example 22 piperidine instead of azetidine in part 6. MS found: (M+H)<sup>+</sup>: 514.

#### Example 29


**4-[(6-Chloroindol-2-yl)sulfonyl]-1-[[4-(iminopyrrolidylmethyl)phenyl]methyl]piperazin-2-one.**



The titled compound was synthesized using a similar procedure to that described in Example 22 pyrrolidine instead of azetidine in part 6. MS found: (M+H)<sup>+</sup>: 500.

#### 20 Example 30


**1-[[4-([2-(Dimethylamino)ethyl]methylamino)iminomethyl]phenyl]methyl]-4-[(6-chloroindol-2-yl)sulfonyl]piperazin-2-one.**



The titled compound was synthesized using a similar procedure to that described in Example 22 using N, N, N'-trimethylethylenediamine instead of azetidine in Part 6. MS found: (M+H)<sup>+</sup>: 531.

### 5 Example 31

**4-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-{[4-(1-methyl(2-imidazolin-2-yl))phenyl]methyl}piperazin-2-one.**




The titled compound was synthesized using a similar procedure to that described in

10 Example 22 using N-methylethylenediamine instead of azetidine in part 6. MS found: (M+H)<sup>+</sup>: 486.

### Example 32

**4-[(5-Chloroindol-2-yl)sulfonyl]-1-({4-**

15 **[imino(methylpropylamino)methyl]phenyl)methyl}piperazin-2-one.**




The titled compound was synthesized using a similar procedure to that described in

Example 22 using N-methyl-N-propylamine instead of azetidine in Part 6. MS found: (M+H)<sup>+</sup>: 502.

20

### Example 33

**1-({4-[(4-Bromopiperidyl)iminomethyl]phenyl)methyl}-4-[(5-chloroindol-2-yl)sulfonyl]piperazin-2-one.**






The titled compound was synthesized using a similar procedure to that described in Example 22 using 4-bromopiperidine instead of azetidine in Part 6. MS found:  $(M+H)^+$ : 592.

5 Example 34


**4-[(5-chloroindol-2-yl)sulfonyl]-1-({4-[imino(methylprop-2-enylamino)methyl]phenyl)methyl}piperazin-2-one.**



The titled compound was synthesized using a similar procedure to that described in Example 22 using N-methylallylamine instead of azetidine in Part 6. MS found:  $(M+H)^+$ : 500.

Example 35

**4-[(5-Chloroindol-2-yl)sulfonyl]-1-[(4-[(2-furylmethyl)methylamino]iminomethyl)phenyl)methyl]piperazin-2-one.**




The titled compound was synthesized using a similar procedure to that described in Example 22 using N-methylfurfurylamine instead of azetidine in Part 6. MS found:  $(M+H)^+$ : 540.

20

Example 36


**1-({4-[(Butylmethylamino)iminomethyl]phenyl)methyl}-4-[(5-chloroindol-2-yl)sulfonyl]piperazin-2-one.**



The titled compound was synthesized using a similar procedure to that described in Example 22 using N-methylbutylamine instead of azetidine in Part 6. MS found:  $(M+H)^+$ : 516.

### 5 Example 37


**4-[(5-Chloroindol-2-yl)sulfonyl]-1-({4-[imino(4-oxopiperidyl)methyl]phenyl}methyl)piperazin-2-one.**



1-({4-[(8-Aza-1,4-dioxaspiro[4.5]dec-8-yl)iminomethyl]phenyl}methyl)-4-[(5-chloroindol-2-yl)sulfonyl]piperazin-2-one from example 24 (10 mg) was dissolved in a solution of 4N HCl in dioxane (3 mL). The mixture was refluxed for 6 hours and concentrated in vacuo. The residue was purified by HPLC (C18 reversed phase) eluting with 0.1% TFA in H<sub>2</sub>O/CH<sub>3</sub>CN to HPLC to afford 4-[(5-Chloroindol-2-yl)sulfonyl]-1-({4-[imino(4-oxopiperidyl)methyl]phenyl}methyl)piperazin-2-one. (7 mg, 71%). MS found:  $(M+H)^+$ : 528.

### Example 38


**4-[(5-Chloroindol-2-yl)sulfonyl]-1-({4-[(cyclohexylmethylamino)iminomethyl]phenyl}methyl)piperazin-2-one.**



The titled compound was synthesized using a similar procedure to that described in Example 22 using N-methylcyclohexylamine instead of azetidine in Part 6. MS found:  $(M+H)^+$ : 542.

### 25 Example 39


**1-{{4-[(5-Chloroindol-2-yl)sulfonyl]-2-oxopiperazinyl}methyl}phenyl}iminomethyl}piperidine-4-carboxylic acid.**



The titled compound was synthesized using a similar procedure to that described in Example 3 using Ethyl 1-{{4-({4-[(5-chloroindol-2-yl)sulfonyl]-2-oxopiperazinyl}methyl)phenyl}iminomethyl}piperidine-4-carboxylate from example 5 23. MS found: (M+H)<sup>+</sup>: 558.

#### Example 40

**4-[(5-Chloroindol-2-yl)sulfonyl]-1-[(4-{imino[methyl(2-phenylethyl)amino]methyl}phenyl)methyl]piperazin-2-one.**




10

The titled compound was synthesized using a similar procedure to that described in Example 22 using N-methylphenethylamine instead of azetidine in Part 6. MS found: (M+H)<sup>+</sup>: 564.

#### 15 Example 41

**4-[(5-Chloroindol-2-yl)sulfonyl]-1-[(4-{imino[methyl(2-(2-pyridyl)ethyl)amino]methyl}phenyl)methyl]piperazin-2-one.**




The titled compound was synthesized using a similar procedure to that described in Example 22 using 4-[2-(methylamino)ethyl]pyridine instead of azetidine in Part 6. MS found: (M+H)<sup>+</sup>: 565.

#### Example 42

**1-({4-[(But-3-ynylmethylamino)iminomethyl]phenyl}methyl)-4-[(5-chloroindol-2-yl)sulfonyl]piperazin-2-one.**

25




The titled compound was synthesized using a similar procedure to that described in Example 22 using N-methyl- -alaninenitrile instead of azetidine in Part 6. MS found:  $(M+H)^+$ : 513.

5

#### Example 43


**4-[(5-Chloroindol-2-yl)sulfonyl]-1-[(4-{imino[methylbenzylamino]methyl}phenyl)methyl]piperazin-2-one.**



10 The titled compound was synthesized using a similar procedure to that described in Example 22 using N-methylbenzylamine instead of azetidine in Part 6. MS found:  $(M+H)^+$ : 550.

#### Example 44


15 **1-[(4-{[(2,2-Dimethoxyethyl)methylamino]iminomethyl}phenyl)methyl]-4-[(5-chloroindol-2-yl)sulfonyl]piperazin-2-one.**



The titled compound was synthesized using a similar procedure to that described in Example 22 using methylaminoacetaldehyde dimethylacetal instead of azetidine in Part  
20 6. MS found:  $(M+H)^+$ : 548.

#### Example 45

**4-[(5-Chloroindol-2-yl)sulfonyl]-1-({4-[imino(4-pyrrolidinylpiperidyl)methyl]phenyl;methyl}piperazin-2-one.**




The titled compound was synthesized using a similar procedure to that described in Example 22 using 4-pyrrolidinopiperidine instead of azetidine in Part 6. MS found: (M+H)<sup>+</sup>: 583.

5

#### Example 46


4-[(5-Chloroindol-2-yl)sulfonyl]-1-[(4-[(1,3-dioxolan-2-ylmethyl)methylamino]iminomethyl}phenyl)methyl]piperazin-2-one.



- 10 The titled compound was synthesized using a similar procedure to that described in Example 22 using 2-methylaminomethyl-1,3-dioxalone instead of azetidine in Part 6. MS found: (M+H)<sup>+</sup>: 546.

#### Example 47

- 15 4-[(5-Chloroindol-2-yl)sulfonyl]-1-[(4-[imino(methylamino)methyl]phenyl)methyl]piperazin-2-one.




The titled compound was synthesized using a similar procedure to that described in Example 22 using methylamine instead of azetidine in Part 6. MS found: (M+H)<sup>+</sup>: 460.

20

#### Example 48

4-[(5-Chloroindol-2-yl)sulfonyl]-1-[(4-{imino[methyl(2-oxoethyl)amino]methyl}phenyl)methyl]piperazin-2-one.




The titled compound was synthesized using a similar procedure to that described in Example 37 using 4-[(5-Chloroindol-2-yl)sulfonyl]-1-[(4-[(1,3-dioxolan-2-ylmethyl)methylamino]iminomethyl]phenyl)methyl]piperazin-2-one from example 46.

5 MS found:  $(M+H)^+$ : 502.

#### Example 49

4-[(5-Chloroindol-2-yl)sulfonyl]-1-[(4-[(ethynylmethylamino)iminomethyl]phenyl)methyl]piperazin-2-one.




10

The titled compound was synthesized using a similar procedure to that described in Example 37 as a minor product. MS found:  $(M+H)^+$ : 484.

#### Example 50

15 4-[(5-Chloroindol-2-yl)sulfonyl]-1-[(4-[imino(methylprop-2-ynylamino)methyl]phenyl)methyl]piperazin-2-one.




The titled compound was synthesized using a similar procedure to that described in Example 22 using N-methylpropargylamine instead of azetidine in Part 6. MS found:

20  $(M+H)^+$ : 498.

#### Example 51


1-[(4-(Azetidinylazetidinylidenemethyl)phenyl)methyl]-4-[(5-chloroindol-2-yl)sulfonyl]piperazin-2-one.



The titled compound was synthesized using a similar procedure to that described in Example 22 as a minor product. MS found:  $(M+H)^+$ : 529.

#### 5 Example 52


**4-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-({4-[imino(methylprop-2-ynylamino)methyl]phenyl)methyl}piperazin-2-one.**



- 10 The titled compound was synthesized using a similar procedure to that described in Example 22 using 5-chloro-2-thienyloxyacetic acid (Ewing, W. R., WO 0032590, 2000) instead of tert-butyl 5-chloro-2-(chlorosulfonyl)indolecarboxylate in Part 5 and N-methylpropargylamine instead of azetidine in Part 6. MS found:  $(M+H)^+$ : 515.

#### 15 Example 53


**4-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-[(4-{imino[methyl(2-phenylethyl)amino]methyl}phenyl)methyl]piperazin-2-one.**



- The titled compound was synthesized using a similar procedure to that described in Example 22 using 5-chloro-2-thienyloxyacetic acid instead of tert-butyl 5-chloro-2-(chlorosulfonyl)indolecarboxylate in Part 5 and N-methylphenethylamine instead of azetidine in Part 6. MS found:  $(M+H)^+$ : 581.

#### Example 54


**4-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-[(4-{imino[methyl(2-(2-pyridyl)ethyl)amino]methyl}phenyl)methyl]piperazin-2-one.**



The titled compound was synthesized using a similar procedure to that described in Example 22 using 5-chloro-2-thienyloxyacetic acid instead of tert-butyl 5-chloro-2-(chlorosulfonyl)indolecarboxylate in Part 5 and 4-[2-(methylamino)ethyl]pyridine instead of azetidine in Part 6. MS found: (M+H)<sup>+</sup>: 582.

Example 55


**10 4-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-({4-[imino(methylprop-2-enylamino)methyl]phenyl)methyl}piperazin-2-one.**



The titled compound was synthesized using a similar procedure to that described in Example 22 using 5-chloro-2-thienyloxyacetic acid instead of tert-butyl 5-chloro-2-(chlorosulfonyl)indolecarboxylate in Part 5 and N-methylallylamine instead of azetidine in Part 6. MS found: (M+H)<sup>+</sup>: 517.

Example 56

**20 4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-[(4-{[(1,3-dioxolan-2-yl)methyl]methylamino}iminomethyl}phenyl)methyl]piperazin-2-one.**




The titled compound was synthesized using a similar procedure to that described in Example 22 using 5-chloro-2-thienyloxyacetic acid instead of tert-butyl 5-chloro-2-(chlorosulfonyl)indolecarboxylate in Part 5 and 2-methylaminomethyl-1,3-dioxalane instead of azetidine in Part 6. MS found: (M+H)<sup>+</sup>: 563.1.



Example 57

**Methyl (2R)-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-[(4-{[(1,3-dioxolan-2-yl)methyl)methylamino]iminomethyl}phenyl)methyl]-6-oxopiperazine-2-carboxylate.**



5


Part 1: To a precooled (ice bath) mixture of Z-D-Dap-OH (7.1 g, 29.8 mmol) and anhydrous MeOH (23 mL), was added slowly SOCl<sub>2</sub> (2.2 mL, 30 mmol). The mixture was allowed to warm to room temperature and stirred at this temperature for 18 hours. The solvent was removed *in vacuo*. The product was crystallized to afford 3-amino-N-  
10 [(benzyloxy)carbonyl]-D-alanine methyl ester hydrochloride (7.5 g, 87%).

Part 2: To a stirred mixture of 3-amino-N-[(benzyloxy)carbonyl]-D-alanine methyl ester from Part 1 (free base: 6.2 g, 24.5 mmol), DIEA (3.5g, 26.7mmol), and anhydrous THF (65 mL), was added methyl bromoacetate (3.7 g, 24.5 mmol) in anhydrous THF  
15 (10 ml). The mixture was stirred at rt for 24 hours. The suspension was filtered and the filtrate evaporated. The residue was treated with Et<sub>2</sub>O, filtered, and the solvent evaporated. The oily residue (7.8 g) was dissolved in EtOH (100 mL) and hydrogenated over 10% Pd/C. The catalyst was filtered off, the solvent removed and the product crystallized from EtOH/Et<sub>2</sub>O to afford 2.7 g (70%) of methyl (2R)-6-  
20 oxopiperazine-2-carboxylate as colorless crystals.

Part 3: The titled compound was synthesized using a similar procedure to that described in Example 1 using methyl (2R)-6-oxopiperazine-2-carboxylate instead of ethyl 3-oxopiperazine-2-acetate in Part 1 and 2-methylaminomethyl-1,3-dioxalone instead of  
25 dimethylamine in Part 3. MS found: (M+H)<sup>+</sup>: 621.

Example 58

(2R)-4-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-[(4-[[[(1,3-dioxolan-2-ylmethyl)methylamino]iminomethyl]phenyl)methyl]-6-oxopiperazine-2-carboxylic acid.




5

- The titled compound was synthesized using a similar procedure to that described in Example 3 using Methyl (2R)-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-[(4-[[[(1,3-dioxolan-2-ylmethyl)methylamino]iminomethyl]phenyl)methyl]-6-oxopiperazine-2-carboxylate. MS found: (M+H)<sup>+</sup>: 607.1.

10

Example 59


Methyl (2R)-1-{[4-(azetidinylinomethyl)phenyl]methyl}-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-6-oxopiperazine-2-carboxylate.



- 15 The titled compound was synthesized using a similar procedure to that described in Example 1 using methyl (2R)-6-oxopiperazine-2-carboxylate instead of ethyl 3-oxopiperazine-2-acetate in Part 1 and azetidine instead of dimethylamine in Part 3. MS found: (M+H)<sup>+</sup>: 561.0.

20 Example 60

(2R)-1-{[4-(Azetidinylinomethyl)phenyl]methyl}-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-6-oxopiperazine-2-carboxylic acid.




The titled compound was synthesized using a similar procedure to that described in

- 25 Example 3. MS found: (M+H)<sup>+</sup>: 547.1.

Example 61

Methyl (2R)-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-[[4-(iminopyrrolidinylmethyl)phenyl]methyl]-6-oxopiperazine-2-carboxylate.




5

The titled compound was synthesized using a similar procedure to that described in Example 1 using methyl (2R)-6-oxopiperazine-2-carboxylate instead of ethyl 3-oxopiperazine-2-acetate in Part 1 and pyrrolidine instead of dimethylamine in Part 3. MS found: (M+H)<sup>+</sup>: 575.1.

10

Example 62


(2R)-4-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-[[4-(iminopyrrolidinylmethyl)phenyl]methyl]-6-oxopiperazine-2-carboxylic acid.



15 The titled compound was synthesized using a similar procedure to that described in Example 3. MS found: (M+H)<sup>+</sup>: 561.1.

Example 63

20 Methyl (2R)-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-[[4-(iminopiperidylmethyl)phenyl]methyl]-6-oxopiperazine-2-carboxylate.




The titled compound was synthesized using a similar procedure to that described in Example 1 using methyl (2R)-6-oxopiperazine-2-carboxylate instead of ethyl 3-oxopiperazine-2-acetate in Part 1 and piperidine instead of dimethylamine in Part 3.

25 MS found: (M+H)<sup>+</sup>: 589.1.

Example 64

(2R)-4-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-[[4-(iminopiperidylmethyl)phenyl]methyl]-6-oxopiperazine-2-carboxylic acid.




5

The titled compound was synthesized using a similar procedure to that described in Example 3. MS found: (M+H)<sup>+</sup>: 575.1.

Example 65


10 Methyl (2R)-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-[(4-{[4-(ethoxycarbonyl)piperidyl]iminomethyl}phenyl)methyl]-6-oxopiperazine-2-carboxylate.



The titled compound was synthesized using a similar procedure to that described in Example 1 using methyl (2R)-6-oxopiperazine-2-carboxylate instead of ethyl 3-oxopiperazine-2-acetate in Part 1 and ethyl isonipecotate instead of dimethylamine in Part 3. MS found: (M+H)<sup>+</sup>: 661.1.

Example 66

20 (2R)-1-({4-[(4-Carboxypiperidyl)iminomethyl]phenyl}methyl)-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-6-oxopiperazine-2-carboxylic acid.




The titled compound was synthesized using a similar procedure to that described in Example 3. MS found: (M+H)<sup>+</sup>: 619.1.

25

Example 67

**Methyl (2R)-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-({4-[imino(methylprop-2-enylamino)methyl]phenyl}methyl)-6-oxopiperazine-2-carboxylate.**




5

The titled compound was synthesized using a similar procedure to that described in Example 1 using methyl (2R)-6-oxopiperazine-2-carboxylate instead of ethyl 3-oxopiperazine-2-acetate in Part 1 and N-methylallylamine instead of dimethylamine in Part 3. MS found:  $(M+H)^+$ : 575.1.

10

Example 68


**(2R)-4-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-({4-[imino(methylprop-2-enylamino)methyl]phenyl}methyl)-6-oxopiperazine-2-carboxylic acid.**



15 The titled compound was synthesized using a similar procedure to that described in Example 3. MS found:  $(M+H)^+$ : 561.1.

Example 69

**Methyl (2R)-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-({4-[imino(methylprop-2-ynylamino)methyl]phenyl}methyl)-6-oxopiperazine-2-carboxylate.**




The titled compound was synthesized using a similar procedure to that described in Example 1 using methyl (2R)-6-oxopiperazine-2-carboxylate instead of ethyl 3-

20

oxopiperazine-2-acetate in Part 1 and N-methylpropargylamine instead of dimethylamine in Part 3. MS found:  $(M+H)^+$ : 573.1.

#### Example 70

- 5 (2R)-4-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-({4-[imino(methylprop-2-ynylamino)methyl]phenyl}methyl)-6-oxopiperazine-2-carboxylic acid.




The titled compound was synthesized using a similar procedure to that described in Example 3. MS found:  $(M+H)^+$ : 559.1.

10

#### Example 71


- Methyl (2R)-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-({4-[imino(methylamino)methyl]phenyl}methyl)-6-oxopiperazine-2-carboxylate.



- 15 The titled compound was synthesized using a similar procedure to that described in Example 1 using methyl (2R)-6-oxopiperazine-2-carboxylate instead of ethyl 3-oxopiperazine-2-acetate in Part 1 and N-methylamine instead of dimethylamine in Part 3. MS found:  $(M+H)^+$ : 535.1.

#### 20 Example 72

- (2R)-4-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-({4-[imino(methylamino)methyl]phenyl}methyl)-6-oxopiperazine-2-carboxylic acid.




The titled compound was synthesized using a similar procedure to that described in

- 25 Example 3. MS found:  $(M+H)^+$ : 521.1.

Example 73


**Methyl (2R)-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-[(4-[(2-cyanoethyl)methylamino]iminomethyl}phenyl)methyl]-6-oxopiperazine-2-**  
**5 carboxylate.**



The titled compound was synthesized using a similar procedure to that described in Example 1 using methyl (2R)-6-oxopiperazine-2-carboxylate instead of ethyl 3-oxopiperazine-2-acetate in Part 1 and N-methyl- -alaninenitrile instead of dimethylamine in Part 3. MS found:  $(M+H)^+$ : 586.1.

Example 74


**(2R)-4-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-[(4-[(2-cyanoethyl)methylamino]iminomethyl}phenyl)methyl]-6-oxopiperazine-2-**  
**15 carboxylic acid.**



The titled compound was synthesized using a similar procedure to that described in Example 3. MS found:  $(M+H)^+$ : 574.1.

Example 75


**Methyl (2R)-1-({4-[(dimethylamino)iminomethyl}phenyl)methyl}-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-6-oxopiperazine-2-carboxylate.**



The titled compound was synthesized using a similar procedure to that described in Example 1 using methyl (2R)-6-oxopiperazine-2-carboxylate instead of ethyl 3-oxopiperazine-2-acetate in Part 1. MS found:  $(M+H)^+$ : 549.1.

#### 5 Example 76


**(2R)-1-({4-[(Dimethylamino)iminomethyl]phenyl}methyl)-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-6-oxopiperazine-2-carboxylic acid.**



The titled compound was synthesized using a similar procedure to that described in Example 3. MS found:  $(M+H)^+$ : 535.1.

#### Example 77

**Methyl (2R)-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-[(4-{imino[methyl(2-phenylethyl)amino]methyl}phenyl)methyl]-6-oxopiperazine-2-carboxylate.**




15

The titled compound was synthesized using a similar procedure to that described in Example 1 using methyl (2R)-6-oxopiperazine-2-carboxylate instead of ethyl 3-oxopiperazine-2-acetate in Part 1 and N-methylphenethylamine instead of dimethylamine in Part 3. MS found:  $(M+H)^+$ : 639.1.

20

#### Example 78

**(2R)-4-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-[(4-{imino[methyl(2-phenylethyl)amino]methyl}phenyl)methyl]-6-oxopiperazine-2-carboxylic acid.**






The titled compound was synthesized using a similar procedure to that described in Example 3. MS found:  $(M+H)^+$ : 625.1.

#### Example 79


- 5 **Methyl (2R)-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-[(4-{imino[methyl(2-(2-pyridyl)ethyl)amino]methyl}phenyl)methyl]-6-oxopiperazine-2-carboxylate.**



The titled compound was synthesized using a similar procedure to that described in Example 1 using methyl (2R)-6-oxopiperazine-2-carboxylate instead of ethyl 3-oxopiperazine-2-acetate in Part 1 and 4-[2-(methylamino)ethyl]pyridine instead of dimethylamine in Part 3. MS found:  $(M+H)^+$ : 640.1.

#### Example 80


- 15 **(2R)-4-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-[(4-{imino[methyl(2-(2-pyridyl)ethyl)amino]methyl}phenyl)methyl]-6-oxopiperazine-2-carboxylic acid.**



The titled compound was synthesized using a similar procedure to that described in Example 3. MS found:  $(M+H)^+$ : 626.1.

#### 20 Example 81

- Methyl (2R)-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-{[4-(1-methyl(2-imidazolin-2-yl))phenyl]methyl}-6-oxopiperazine-2-carboxylate.**




The titled compound was synthesized using a similar procedure to that described in Example 1 using methyl (2R)-6-oxopiperazine-2-carboxylate instead of ethyl 3-oxopiperazine-2-acetate in Part 1 and 4-[2-(methylamino)ethyl]pyridine instead of dimethylamine in Part 3. MS found:  $(M+H)^+$ : 640.1.

oxopiperazine-2-acetate in Part 1 and N-methylethylenediamine instead of dimethylamine in Part 3. MS found:  $(M+H)^+$ : 561.

### Example 82

- 5 (2R)-4-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-{[4-(1-methyl(2-imidazolin-2-yl))phenyl]methyl}-6-oxopiperazine-2-carboxylic acid.




The titled compound was synthesized using a similar procedure to that described in Example 3. MS found:  $(M+H)^+$ : 547.

10

### Example 83

- Methyl (2R)-1-[(4-{[(1,3-dioxolan-2-ylmethyl)methylamino]iminomethyl}phenyl)methyl]-4-[(5-methylindol-2-yl)sulfonyl]-6-oxopiperazine-2-carboxylate.


15



The titled compound was synthesized using a similar procedure to that described in Example 22. MS found:  $(M+H)^+$ : 604.

### 20 Example 84


- (2R)-4-[(5-Chloroindol-2-yl)sulfonyl]-1-[(4-{[(1,3-dioxolan-2-ylmethyl)methylamino]iminomethyl}phenyl)methyl]-6-oxopiperazine-2-carboxylic acid.



The titled compound was synthesized using a similar procedure to that described in Example 3. MS found:  $(M+H)^+$ : 590.

Example 85

- 5 **Methyl (2R)-1-[[4-(azetidinyliminomethyl)phenyl]methyl]-4-[(5-chloroindol-2-yl)sulfonyl]-6-oxopiperazine-2-carboxylate.**




The titled compound was synthesized using a similar procedure to that described in Example 22. MS found:  $(M+H)^+$ : 544.

10

Example 86

- (2R)-1-[[4-(Azetidinyliminomethyl)phenyl]methyl]-4-[(5-chloroindol-2-yl)sulfonyl]-6-oxopiperazine-2-carboxylic acid.




15

The titled compound was synthesized using a similar procedure to that described in Example 3. MS found:  $(M+H)^+$ : 530.1.

Example 87

- 20 **Methyl (2R)-4-[(5-chloroindol-2-yl)sulfonyl]-1-[[4-(iminopyrrolidinymethyl)phenyl]methyl]-6-oxopiperazine-2-carboxylate.**




The titled compound was synthesized using a similar procedure to that described in Example 22. MS found:  $(M+H)^+$ : 558.

25

Example 88

(2R)-4-[(5-Chloroindol-2-yl)sulfonyl]-1-[[4-(iminopyrrolidinylmethyl)phenyl]methyl]-6-oxopiperazine-2-carboxylic acid.




- 5 The titled compound was synthesized using a similar procedure to that described in Example 3. MS found:  $(M+H)^+$ : 544.

Example 89

Methyl (2R)-4-[(5-chloroindol-2-yl)sulfonyl]-1-[[4-


- 10 (iminopiperidylmethyl)phenyl]methyl]-6-oxopiperazine-2-carboxylate.



The titled compound was synthesized using a similar procedure to that described in Example 22. MS found:  $(M+H)^+$ : 572.

15 Example 90

(2R)-4-[(5-Chloroindol-2-yl)sulfonyl]-1-[[4-(iminopiperidylmethyl)phenyl]methyl]-6-oxopiperazine-2-carboxylic acid.



The titled compound was synthesized using a similar procedure to that described in


- 20 Example 3. MS found:  $(M+H)^+$ : 558.

Example 91

Methyl (2R)-4-[(5-chloroindol-2-yl)sulfonyl]-1-[(4-[[4-(ethoxycarbonyl)piperidyl]iminomethyl]phenyl)methyl]-6-oxopiperazine-2-

25 carboxylate.


84



The titled compound was synthesized using a similar procedure to that described in Example 22. MS found:  $(M+H)^+$ : 644.

#### 5 Example 92

**(2R)-1-({4-[(4-Carboxypiperidyl)iminomethyl]phenyl}methyl)-4-[(5-chloroindol-2-yl)sulfonyl]-6-oxopiperazine-2-carboxylic acid.**




The titled compound was synthesized using a similar procedure to that described in

10 Example 3. MS found:  $(M+H)^+$ : 602.

#### Example 93

**Methyl (2R)-4-[(5-chloroindol-2-yl)sulfonyl]-1-({4-[imino(methylprop-2-enylamino)methyl]phenyl}methyl)-6-oxopiperazine-2-carboxylate.**




15

The titled compound was synthesized using a similar procedure to that described in Example 22. MS found:  $(M+H)^+$ : 558.

#### Example 94


20 **(2R)-4-[(5-Chloroindol-2-yl)sulfonyl]-1-({4-[imino(methylprop-2-enylamino)methyl]phenyl}methyl)-6-oxopiperazine-2-carboxylic acid.**



The titled compound was synthesized using a similar procedure to that described in Example 3. MS found:  $(M+H)^+$ : 544.0.

Example 95

- 5 **Methyl (2R)-4-[(5-chloroindol-2-yl)sulfonyl]-1-({4-[imino(methylprop-2-ynylamino)methyl]phenyl}methyl)-6-oxopiperazine-2-carboxylate.**




The titled compound was synthesized using a similar procedure to that described in Example 22. MS found:  $(M+H)^+$ : 556.1.

10

Example 96


- (2R)-4-[(5-Chloroindol-2-yl)sulfonyl]-1-({4-[imino(methylprop-2-ynylamino)methyl]phenyl}methyl)-6-oxopiperazine-2-carboxylic acid.



- 15 The titled compound was synthesized using a similar procedure to that described in Example 3. MS found:  $(M+H)^+$ : 542.

Example 97


- Methyl (2R)-4-[(5-chloroindol-2-yl)sulfonyl]-1-({4-[imino(methylamino)methyl]phenyl}methyl)-6-oxopiperazine-2-carboxylate.**



The titled compound was synthesized using a similar procedure to that described in Example 22. MS found:  $(M+H)^+$ : 518.1.

- 25 Example 98


(2R)-4-[(5-Chloroindol-2-yl)sulfonyl]-1-({4-[imino(methylamino)methyl]phenyl)methyl}-6-oxopiperazine-2-carboxylic acid.



The titled compound was synthesized using a similar procedure to that described in  
 5 Example 3. MS found:  $(M+H)^+$ : 504.

#### Example 99


Methyl (2R)-4-[(5-chloroindol-2-yl)sulfonyl]-1-[(4-[(2-cyanoethyl)methylamino]iminomethyl]phenyl)methyl]-6-oxopiperazine-2-  
 10 carboxylate.



The titled compound was synthesized using a similar procedure to that described in  
 Example 22. MS found:  $(M+H)^+$ : 571.

#### 15 Example 100

(2R)-4-[(5-Chloroindol-2-yl)sulfonyl]-1-[(4-[(2-cyanoethyl)methylamino]iminomethyl]phenyl)methyl]-6-oxopiperazine-2-carboxylic acid.




20 The titled compound was synthesized using a similar procedure to that described in  
 Example 3. MS found:  $(M+H)^+$ : 557.1.

#### Example 101

Methyl (2R)-1-({4-[(dimethylamino)iminomethyl]phenyl)methyl}-4-[(5-  
 25 chloroindol-2-yl)sulfonyl]-6-oxopiperazine-2-carboxylate.


87



The titled compound was synthesized using a similar procedure to that described in Example 22. MS found:  $(M+H)^+$ : 532.1.

#### 5 Example 102

**(2R)-1-({4-[(Dimethylamino)iminomethyl]phenyl}methyl)-4-[(5-chloroindol-2-yl)sulfonyl]-6-oxopiperazine-2-carboxylic acid.**




The titled compound was synthesized using a similar procedure to that described in

10 Example 3. MS found:  $(M+H)^+$ : 518.1.

#### Example 103

**Methyl (2R)-4-[(5-chloroindol-2-yl)sulfonyl]-1-[(4-{imino[methyl(2-phenylethyl)amino]methyl}phenyl)methyl]-6-oxopiperazine-2-carboxylate.**




15

The titled compound was synthesized using a similar procedure to that described in Example 22. MS found:  $(M+H)^+$ : 622.1.

#### Example 104

20 **(2R)-4-[(5-Chloroindol-2-yl)sulfonyl]-1-[(4-{imino[methyl(2-phenylethyl)amino]methyl}phenyl)methyl]-6-oxopiperazine-2-carboxylic acid.**






The titled compound was synthesized using a similar procedure to that described in Example 3. MS found:  $(M+H)^+$ : 608.1.

#### Example 105

- 5 Methyl (2R)-4-[(5-chloroindol-2-yl)sulfonyl]-1-[(4-{imino[methyl(2-(2-pyridyl)ethyl)amino]methyl}phenyl)methyl]-6-oxopiperazine-2-carboxylate.




The titled compound was synthesized using a similar procedure to that described in Example 22. MS found:  $(M+H)^+$ : 623.1.

10

#### Example 106


- (2R)-4-[(5-Chloroindol-2-yl)sulfonyl]-1-[(4-{imino[methyl(2-(2-pyridyl)ethyl)amino]methyl}phenyl)methyl]-6-oxopiperazine-2-carboxylic acid.



- 15 The titled compound was synthesized using a similar procedure to that described in Example 3. MS found:  $(M+H)^+$ : 609.1.

#### Example 107


- 20 Methyl (2R)-4-[(5-chloroindol-2-yl)sulfonyl]-1-[[4-(1-methyl(2-imidazolin-2-yl))phenyl)methyl]-6-oxopiperazine-2-carboxylate.



The titled compound was synthesized using a similar procedure to that described in Example 22. MS found:  $(M+H)^+$ : 544.1.

Example 108


(2R)-4-[(5-Chloroindol-2-yl)sulfonyl]-1-[[4-(1-methyl(2-imidazolin-2-yl))phenyl)methyl]-6-oxopiperazine-2-carboxylic acid.



- 5 The titled compound was synthesized using a similar procedure to that described in Example 3. MS found:  $(M+H)^+$ : 530.

Example 109


Methyl (2R)-4-[(5-chloroindol-2-yl)sulfonyl]-1-[(4-{imino[methyl(2-oxoethyl)amino]methyl}phenyl)methyl]-6-oxopiperazine-2-carboxylate.



The titled compound was synthesized using a similar procedure to that described in Example 37 using compound of example 83. MS found:  $(M+H)^+$ : 560.1.

15 Example 110

(2R)-4-[(5-Chloroindol-2-yl)sulfonyl]-1-[(4-{imino[methyl(2-oxoethyl)amino]methyl}phenyl)methyl]-6-oxopiperazine-2-carboxylic acid.




- The titled compound was synthesized using a similar procedure to that described in Example 3. MS found:  $(M+H)^+$ : 546.1.

Example 111

Methyl (2R)-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-[(4-{imino[methyl(2-oxoethyl)amino]methyl}phenyl)methyl]-6-oxopiperazine-2-carboxylate.


90



The titled compound was synthesized using a similar procedure to that described in Example 37 using compound of example 57. MS found:  $(M+H)^+$ : 577.

#### 5 Example 112

**(2R)-4-[(6-Chlorobenzo[b]thiophen-2-yl)sulfonyl]-1-[(4-{imino[methyl(2-oxoethyl)amino]methyl}phenyl)methyl]-6-oxopiperazine-2-carboxylic acid.**




The titled compound was synthesized using a similar procedure to that described in

10 Example 3. MS found:  $(M+H)^+$ : 563.

#### Example 113

**Methyl (2R)-1-[[4-(azetidinylazetidinyldenemethyl)phenyl]methyl]-4-[(5-chloroindol-2-yl)sulfonyl]-6-oxopiperazine-2-carboxylate.**




15

The titled compound was synthesized using a similar procedure to that described in Example 22 as a minor product in the synthesis of compound of example 85. MS found:  $(M+H)^+$ : 585.

#### 20 Example 114


**(2R)-1-[[4-(Azetidinylazetidinyldenemethyl)phenyl]methyl]-4-[(5-chloroindol-2-yl)sulfonyl]-6-oxopiperazine-2-carboxylic acid.**



The titled compound was synthesized using a similar procedure to that described in Example 3. MS found:  $(M+H)^+$ : 571.

#### 5 Example 115


**Methyl (2R)-1-{{4-(azetidinylazetidinylidenemethyl)phenyl}methyl}-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-6-oxopiperazine-2-carboxylate.**



The titled compound was synthesized using a similar procedure to that described in  
 10 Example 1 in the synthesis of compound of example 59 as a minor product. MS found:  $(M+H)^+$ : 602.

#### Example 116

**(2R)-1-{{4-(Azetidinylazetidinylidenemethyl)phenyl}methyl}-4-[(6-chlorobenzo[b]thiophen-2-yl)sulfonyl]-6-oxopiperazine-2-carboxylic acid.**



The titled compound was synthesized using a similar procedure to that described in Example 3. MS found:  $(M+H)^+$ : 588.

#### 20 Compositions and Formulations

Compositions or formulations of the compounds of the invention are prepared for storage or administration by mixing the compound having a desired degree of purity with physiologically acceptable carriers, excipients, stabilizers etc., and may be

provided in sustained release or timed release formulations. Acceptable carriers or diluents for therapeutic use are well known in the pharmaceutical field, and are described, for example, in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., (A.R. Gennaro edit. 1985). Such materials are nontoxic to the recipients at the  
5 dosages and concentrations employed, and include buffers such as phosphate, citrate, acetate and other organic acid salts, antioxidants such as ascorbic acid, low molecular weight (less than about ten residues) peptides such as polyarginine, proteins, such as serum albumin, gelatin, or immunoglobulins, hydrophilic polymers such as polyvinylpyrrolidinone, amino acids such as glycine, glutamic acid, aspartic acid, or  
10 arginine, monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, mannose or dextrans, chelating agents such as EDTA, sugar alcohols such as mannitol or sorbitol, counterions such as sodium and/or nonionic surfactants such as Tween®, Pluronic® or polyethyleneglycol.

Dosage formulations of the compounds of the invention to be used for  
15 therapeutic administration must be sterile. Sterility is readily accomplished by filtration through sterile membranes such as 0.2 micron membranes, or by other conventional methods. Formulations typically will be stored in lyophilized form or as an aqueous solution. The pH of the preparations of the invention typically will be between about 3 and about 11, more preferably from about 5 to about 9 and most preferably from about  
20 7 to about 8. It will be understood that use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of cyclic polypeptide salts. While the preferred route of administration is by injection, other methods of administration are also anticipated such as intravenously (bolus and/or infusion), subcutaneously, intramuscularly, colonically, rectally, nasally or intraperitoneally, employing a variety  
25 of dosage forms such as suppositories, implanted pellets or small cylinders, aerosols, oral dosage formulations and topical formulations such as ointments, drops and dermal patches. The compounds of the invention are desirably incorporated into shaped articles such as implants which may employ inert materials such as biodegradable polymers or synthetic silicones, for example, Silastic, silicone rubber or other polymers  
30 commercially available.

The compounds of the invention may also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of lipids, such as cholesterol, stearylamine or phosphatidylcholines.

5 The compounds of the invention may also be delivered by the use of antibodies, antibody fragments, growth factors, hormones, or other targeting moieties, to which the compound molecules are coupled. The compounds of the invention may also be coupled with suitable polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxy-propyl-methacrylamide-phenol, 10 polyhydroxyethyl-aspartamide-phenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, the factor Xa inhibitors of the invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, 15 polyacetals, polydihydropyrans, polycyanoacrylates and cross linked or amphipathic block copolymers of hydrogels. Polymers and semipermeable polymer matrices may be formed into shaped articles, such as valves, stents, tubing, prostheses and the like.

Therapeutic compound liquid formulations generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a 20 stopper pierceable by hypodermic injection needle.

Therapeutically effective dosages may be determined by either *in vitro* or *in vivo* methods. For each particular compound of the present invention, individual determinations may be made to determine the optimal dosage required. The range of therapeutically effective dosages will naturally be influenced by the route of 25 administration, the therapeutic objectives, and the condition of the patient. For injection by hypodermic needle, it may be assumed the dosage is delivered into the body's fluids. For other routes of administration, the absorption efficiency must be individually determined for each inhibitor by methods well known in pharmacology. Accordingly, it may be necessary for the therapist to titer the dosage and modify the 30 route of administration as required to obtain the optimal therapeutic effect. The determination of effective dosage levels, that is, the dosage levels necessary to achieve

the desired result, will be within the ambit of one skilled in the art. Typically, applications of compound are commenced at lower dosage levels, with dosage levels being increased until the desired effect is achieved.

A typical dosage of the compounds and compositions of the invention range  
5 from about 0.001 mg/kg to about 1000 mg/kg, preferably from about 0.01 mg/kg to about 100 mg/kg, and more preferably from about 0.10 mg/kg to about 20 mg/kg. Advantageously, the compounds of the invention may be administered several times daily. Other dosage regimens may also be useful (e.g. single daily dose and/or continuous infusion).

10 Typically, about 0.5 to about 500 mg of a compound or mixture of compounds of the invention, as the free acid or base form or as a pharmaceutically acceptable salt, is compounded with a physiologically acceptable vehicle, carrier, excipient, binder, preservative, stabilizer, dye, flavor, etc., as called for by accepted pharmaceutical practice. The amount of active ingredient in these compositions is such that a suitable  
15 dosage in the range indicated is obtained.

Typical adjuvants which may be incorporated into tablets, capsules and the like are a binder such as acacia, corn starch or gelatin, and excipient such as microcrystalline cellulose, a disintegrating agent like corn starch or alginic acid, a lubricant such as magnesium stearate, a sweetening agent such as sucrose or lactose, or  
20 a flavoring agent. When a dosage form is a capsule, in addition to the above materials it may also contain a liquid carrier such as water, saline, a fatty oil. Other materials of various types may be used as coatings or as modifiers of the physical form of the dosage unit. Sterile compositions for injection can be formulated according to conventional pharmaceutical practice. For example, dissolution or suspension of the  
25 active compound in a vehicle such as an oil or a synthetic fatty vehicle like ethyl oleate, or into a liposome may be desired. Buffers, preservatives, antioxidants and the like can be incorporated according to accepted pharmaceutical practice.

The preferred compounds of the present invention are characterized by their ability to inhibit thrombus formation with acceptable effects on classical measures of  
30 coagulation parameters, platelets and platelet function, and acceptable levels of bleeding complications associated with their use. Conditions characterized by

undesired thrombosis would include those involving the arterial and venous vasculature.

With respect to the coronary arterial vasculature, abnormal thrombus formation characterizes the rupture of an established atherosclerotic plaque which is the major  
5 cause of acute myocardial infarction and unstable angina, as well as also characterizing the occlusive coronary thrombus formation resulting from either thrombolytic therapy or percutaneous transluminal coronary angioplasty (PTCA).

With respect to the venous vasculature, abnormal thrombus formation characterizes the condition observed in patients undergoing major surgery in the lower  
10 extremities or the abdominal area who often suffer from thrombus formation in the venous vasculature resulting in reduced blood flow to the affected extremity and a predisposition to pulmonary embolism. Abnormal thrombus formation further characterizes disseminated intravascular coagulopathy commonly occurs within both vascular systems during septic shock, certain viral infections and cancer, a condition  
15 wherein there is rapid consumption of coagulation factors and systemic coagulation which results in the formation of life-threatening thrombi occurring throughout the microvasculature leading to widespread organ failure.

The compounds of the invention are useful for the treatment or prophylaxis of those diseases which involve the production and/or action of factor Xa/prothrombinase  
20 complex. The compounds of this present invention, selected and used as disclosed herein, find utility as a diagnostic or therapeutic agent for preventing or treating a condition in a mammal characterized by undesired thrombosis or a disorder of coagulation. Disease states treatable or preventable by the administration of compounds of the invention include, without limitation, occlusive coronary thrombus  
25 formation resulting from either thrombolytic therapy or percutaneous transluminal coronary angioplasty, thrombus formation in the venous vasculature, disseminated intravascular coagulopathy, the treatment of reocclusion or restenosis of reperfused coronary arteries, thromboembolic complications of surgery and peripheral arterial occlusion, a condition wherein there is rapid consumption of coagulation factors and  
30 systemic coagulation which results in the formation of life-threatening thrombi occurring throughout the microvasculature leading to widespread organ failure,



hemorrhagic stroke, renal dialysis, blood oxygenation, and cardiac catheterization.

Accordingly, the invention provides a method for preventing or treating a condition in a mammal characterized by undesired thrombosis which administers to a mammal a therapeutically effective amount of a compound of the invention, as

5 described herein. Conditions for prevention or treatment include, for example, (a) the treatment or prevention of any thrombotically mediated acute coronary syndrome including myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, (b) the treatment or prevention of any thrombotically mediated cerebrovascular syndrome

10 including embolic stroke, thrombotic stroke or transient ischemic attacks, (c) the treatment or prevention of any thrombotic syndrome occurring in the venous system including deep venous thrombosis or pulmonary embolus occurring either spontaneously or in the setting of malignancy, surgery or trauma, (d) the treatment or prevention of any coagulopathy including disseminated intravascular coagulation

15 (including the setting of septic shock or other infection, surgery, pregnancy, trauma or malignancy and whether associated with multi-organ failure or not), thrombotic thrombocytopenic purpura, thromboangiitis obliterans, or thrombotic disease associated with heparin induced thrombocytopenia, (e) the treatment or prevention of thrombotic complications associated with extracorporeal circulation (e.g. renal dialysis,

20 cardiopulmonary bypass or other oxygenation procedure, plasmapheresis), (f) the treatment or prevention of thrombotic complications associated with instrumentation (e.g. cardiac or other intravascular catheterization, intra-aortic balloon pump, coronary stent or cardiac valve), and (g) those involved with the fitting of prosthetic devices.

Anticoagulant therapy is also useful to prevent coagulation of stored whole

25 blood and to prevent coagulation in other biological samples for testing or storage.

Thus the compounds of the invention can be added to or contacted with any medium containing or suspected to contain factor Xa and in which it is desired that blood coagulation be inhibited, e.g., when contacting the mammal's blood with material such as vascular grafts, stents, orthopedic prostheses, cardiac stents, valves and prostheses,

30 extra corporeal circulation systems and the like.

Thus, the compounds of the invention also find utility in a method for inhibiting the coagulation of biological samples by administration of a compound of the invention.

5

### BIOLOGICAL ACTIVITY EXAMPLES

Evaluation of the compounds of the invention is guided by in vitro protease activity assays (see below) and in vivo studies to evaluate antithrombotic efficacy, and effects on hemostasis and hematological parameters.

The compounds of the present invention are dissolved in buffer to give solutions containing concentrations such that assay concentrations range from about 0 to 100  $\mu\text{M}$ . In the assays for thrombin, prothrombinase and factor Xa, a synthetic chromogenic substrate is added to a solution containing test compound and the enzyme of interest and the residual catalytic activity of that enzyme is determined spectrophotometrically. The  $\text{IC}_{50}$  of a compound is determined from the substrate turnover. The  $\text{IC}_{50}$  is the concentration of test compound giving 50% inhibition of the substrate turnover. The compounds of the present invention desirably have an  $\text{IC}_{50}$  of less than about 500 nM in the factor Xa assay, preferably less than about 200 nM, and more preferred compounds have an  $\text{IC}_{50}$  of about 100 nM or less in the factor Xa assay. The compounds of the present invention desirably have an  $\text{IC}_{50}$  of less than about 4.0  $\mu\text{M}$  in the prothrombinase assay, preferably less than 200 nM, and more preferred compounds have an  $\text{IC}_{50}$  of about 10 nM or less in the prothrombinase assay. The compounds of the present invention desirably have an  $\text{IC}_{50}$  of greater than about 1.0  $\mu\text{M}$  in the thrombin assay, preferably greater than about 10.0  $\mu\text{M}$ , and more preferred compounds have an  $\text{IC}_{50}$  of greater than about 100.0  $\mu\text{M}$  in the thrombin assay.

#### Amidolytic Assays for determining protease inhibition activity

The factor Xa and thrombin assays are performed at room temperature, in 0.02 M Tris-HCl buffer, pH 7.5, containing 0.15 M NaCl. The rates of hydrolysis of the para-nitroanilide substrate S-2765 (Chromogenix) for factor Xa, and the substrate

Chromozym TH (Boehringer Mannheim) for thrombin following preincubation of the enzyme with inhibitor for 5 minutes at room temperature, and were determined using the Softmax 96-well plate reader (Molecular Devices), monitored at 405 nm to measure the time dependent appearance of p-nitroaniline.

5       The prothrombinase inhibition assay is performed in a plasma free system with modifications to the method described by Sinha, U. *et al.*, *Thromb. Res.*, 75, 427-436 (1994). Specifically, the activity of the prothrombinase complex is determined by measuring the time course of thrombin generation using the p-nitroanilide substrate Chromozym TH. The assay consists of preincubation ( 5 minutes) of selected  
10 compounds to be tested as inhibitors with the complex formed from factor Xa (0.5 nM), factor Va (2 nM), phosphatidyl serine:phosphatidyl choline (25:75, 20  $\mu$ M) in 20 mM Tris-HCl buffer, pH 7.5, containing 0.15 M NaCl, 5 mM CaCl<sub>2</sub> and 0.1% bovine serum albumin. Aliquots from the complex-inhibitor mixture are added to prothrombin (1 nM) and Chromozym TH (0.1 mM). The rate of substrate cleavage is monitored at 405  
15 nm for two minutes. Eight different concentrations of inhibitor are assayed in duplicate. A standard curve of thrombin generation by an equivalent amount of untreated complex are used for determination of percent inhibition.

#### Antithrombotic Efficacy in a Rabbit Model of Venous Thrombosis

20       A rabbit deep vein thrombosis model as described by Hollenbach, S. et al., *Thromb. Haemost.* 71, 357-362 (1994), is used to determine the in-vivo antithrombotic activity of the test compounds. Rabbits are anesthetized with I.M. injections of Ketamine, Xylazine, and Acepromazine cocktail. A standardized protocol consists of insertion of a thrombogenic cotton thread and copper wire apparatus into the abdominal  
25 vena cava of the anesthetized rabbit. A non-occlusive thrombus is allowed to develop in the central venous circulation and inhibition of thrombus growth is used as a measure of the antithrombotic activity of the studied compounds. Test agents or control saline are administered through a marginal ear vein catheter. A femoral vein catheter is used for blood sampling prior to and during steady state infusion of test compound.  
30 Initiation of thrombus formation begins immediately after advancement of the cotton thread apparatus into the central venous circulation. Test compounds are administered

from time = 30 min to time = 150 min at which the experiment is terminated. The rabbits are euthanized and the thrombus excised by surgical dissection and characterized by weight and histology. Blood samples are analyzed for changes in hematological and coagulation parameters.

5


Effects of Compounds in Rabbit Venous Thrombosis model

Administration of compounds in the rabbit venous thrombosis model demonstrates antithrombotic efficacy at the higher doses evaluated. There are no significant effects of the compound on the aPTT and PT prolongation with the highest  
10 dose (100  $\mu\text{g/kg}$  + 2.57  $\mu\text{g/kg/min}$ ). Compounds have no significant effects on hematological parameters as compared to saline controls. All measurements are an average of all samples after steady state administration of vehicle or (D)-Arg-Gly-Arg-thiazole. Values are expressed as mean  $\pm$  SD.

15 Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and the following illustrative examples, make and utilize the compounds of the present invention and practice the claimed methods. It should be understood that the foregoing discussion and examples merely present a detailed description of certain preferred embodiments. It will be apparent to those of  
20 ordinary skill in the art that various modifications and equivalents can be made without departing from the spirit and scope of the invention. All the patents, journal articles and other documents discussed or cited above are herein incorporated by reference.


WHAT IS CLAIMED IS:

1. A compound of the formulae (I) or (II):



wherein:

- 5 A is a member selected from the group consisting of:



- $R^{1a}$ ,  $R^{1b}$ ,  $R^{1d}$ , and  $R^{1e}$  are each independently a H,  $C_{1-6}$  alkyl,  $C_{2-6}$  alkenyl,  $C_{2-6}$  alkynyl,  $C_{3-8}$  cycloalkyl, aryl,  $-C_{1-6}$  alkylaryl, heterocyclyl,  $-C_{1-6}$  alkylheterocyclyl,  $-(CH_2)_{1-6}OH$ ,   
 10  $-(CH_2)_{1-6}OC_{1-6}$  alkyl,  $-(CH_2)_{1-6}NH_2$ ,  $-(CH_2)_{1-6}NHC_{1-6}$  alkyl,  $-(CH_2)_{1-6}N(C_{1-6} \text{ alkyl})_2$ ,   
 $-(CH_2)_{1-6}CHNH(COOH)$ ,  $-(CH_2)_{1-6}NHC(=O)C_{1-6}$  alkyl,  $-(CH_2)_{1-6}CHO$ ,  $-(CH_2)_{1-6}C(=O)OH$ ,   
 $-(CH_2)_{1-6}C(=O)OC_{1-6}$  alkyl, or  $-(CH_2)_{1-6}C(=O)NH_2$ ; wherein  $R^{1a}$ ,  $R^{1b}$ ,  $R^{1d}$ , or  $R^{1e}$  is optionally substituted with at least one of halo, alkyl, alkylideneamine, arylideneamine, cyano, hydroxy, alkoxy, amino, amidino, guanidino, imino, amido, acid,   
 15 ester, keto, aldehyde, dioxolane, furanyl, piperidiny, piperaziny, pyrrolidiny, aryl, morpholiny, and thiomorpholinyldioxide; or  $R^{1a}$  and  $R^{1b}$  or  $R^{1a}$  and  $R^{1c}$  or  $R^{1a}$  and  $R^{1d}$  or  $R^{1d}$  and  $R^{1e}$  taken together with the nitrogen atom to which they are each attached can form a substituted or unsubstituted 3 to 8 membered heterocyclic or heteroaromatic amine group which, optionally, contains at least one other heteroatom of N, O or S;

wherein  $R^{1a}$ ,  $R^{1b}$ ,  $R^{1d}$ , or  $R^{1e}$  is optionally substituted with at least one of halo, alkyl, alkylideneamine, arylideneamine, cyano, hydroxy, alkoxy, amino, amidino, guanidino, imino, amido, acid, ester, keto, aldehyde, dioxolane, furanyl, piperidinyl, piperazinyl, pyrrolidinyl, aryl, morpholinyl, and thiomorpholinyldioxide;

5

$R^{1c}$  is H,  $C_{1-6}$ alkyl or  $C_{3-8}$ cycloalkyl;

$R^{2a}$ ,  $R^{2b}$  and  $R^{2c}$  are each independently a H,  $C_{1-6}$ alkyl,  $C_{2-6}$ alkenyl,  $C_{2-6}$ alkynyl,  $C_{3-8}$ cycloalkyl, aryl,  $-C_{1-6}$ alkylaryl, heterocyclyl,  $-C_{1-6}$ alkylheterocyclyl,  $-(CH_2)_{1-6}OH$ ,  $-(CH_2)_{1-6}OC_{1-6}$ alkyl,  $-(CH_2)_{1-6}NH_2$ ,  $-(CH_2)_{1-6}NHC_{1-6}$ alkyl,  $-(CH_2)_{1-6}N(C_{1-6}alkyl)_2$ ,  $-(CH_2)_{1-6}CHNH(COOH)$ ,  $-(CH_2)_{1-6}NHC(=O)C_{1-6}$ alkyl,  $-(CH_2)_{1-6}CHO$ ,  $-(CH_2)_{1-6}C(=O)OH$ ,  $-(CH_2)_{1-6}C(=O)OC_{1-6}$ alkyl, or  $-(CH_2)_{1-6}C(=O)NH_2$ ; wherein  $R^{2a}$ ,  $R^{2b}$  and  $R^{2c}$  is optionally substituted with at least one of halo, alkyl, alkylideneamine, arylideneamine, cyano, hydroxy, alkoxy, amino, amidino, guanidino, imino, amido, acid, ester, keto, aldehyde, dioxolane, furanyl, piperidinyl, piperazinyl, pyrrolidinyl, aryl, morpholinyl, and thiomorpholinyldioxide; or  $R^{2a}$  and  $R^{2b}$  or  $R^{1a}$ , as set forth above, and  $R^{2a}$  or  $R^{1a}$ , as set forth above, and  $R^{2b}$  taken together with the nitrogen atom to which they are each attached can form a substituted or unsubstituted 3 to 8 membered heterocyclic or heteroaromatic amine group which, optionally, contains at least one other heteroatom of N, O or S; wherein  $R^{2a}$ ,  $R^{2b}$  or  $R^{2c}$  is optionally substituted with at least one of halo, alkyl, alkylideneamine, arylideneamine, cyano, hydroxy, alkoxy, amino, amidino, guanidino, imino, amido, acid, ester, keto, aldehyde, dioxolane, furanyl, piperidinyl, piperazinyl, pyrrolidinyl, aryl, morpholinyl, and thiomorpholinyldioxide;


25

R is, in each occurrence, independently, H,  $-C_{1-6}$ alkyl,  $-C_{3-8}$ cycloalkyl,  $-C_{0-6}$ alkyl- $OC_{1-6}$ alkyl,  $-C_{0-6}$ alkyl- $O(CH_2)_{1-4}-COOH$ ,  $-C_{0-6}$ alkyl- $O(CH_2)_{1-4}-C(=O)OC_{1-6}$ alkyl,  $-C_{0-6}$ alkylCOOH,  $-C_{0-6}$ alkylCO<sub>2</sub> $C_{1-6}$ alkyl,  $-C_{0-6}$ alkylOC<sub>1-6</sub>alkyl,  $-C_{1-6}$ alkylOH,  $-C_{0-6}$ alkylCONH<sub>2</sub>,  $-C_{0-6}$ alkylCONHC<sub>0-6</sub>alkyl,  $-C_{0-6}$ alkylCON( $C_{0-6}$ alkyl)<sub>2</sub>,  $-C_{0-6}$ alkylCON( $CH_2$ )<sub>2-6</sub>,  $-C_{0-6}$ alkylCON( $CH_2CH_2$ )<sub>2</sub>O,  $-C_{0-6}$ alkylCON( $CH_2CH_2$ )<sub>2</sub>SO<sub>2</sub>

30

-C<sub>0-6</sub>alkylCONHaryl, -C<sub>0-6</sub>alkylNH<sub>2</sub>, -C<sub>0-6</sub>alkylNH(C<sub>1-6</sub>alkyl) or  
-C<sub>0-6</sub>alkylN(C<sub>1-6</sub>alkyl)<sub>2</sub>.


Q is a member selected from the group consisting of:



Y is S;

R<sup>1</sup> is H, -Cl, -Br, -I, -F, -OCF<sub>3</sub>, alkyl, hydroxy, alkoxy, amino, thiol, thioalkyl, thioaryl,  
10 or piperiziny;

J<sup>1</sup> is a member selected from the group consisting of:




X is O or S;

R<sup>2</sup> is H, -Cl, -Br, -I, -F or -OC<sub>1-6</sub>alkyl;

R<sup>3</sup> is H, -Cl, -Br, -I, -F, -OC<sub>1-6</sub>alkyl, -NHC<sub>1-6</sub>acyl, -NO<sub>2</sub>, -NHSO<sub>2</sub>C<sub>1-4</sub>alkyl, -CN,  
-NH<sub>2</sub>, -CONH<sub>2</sub>, -SO<sub>2</sub>C<sub>1-6</sub>alkyl, -SO<sub>2</sub>NH<sub>2</sub>, -CO<sub>2</sub>C<sub>1-6</sub>alkyl or -O(CH<sub>2</sub>)<sub>1-4</sub>COOH;

20 R<sup>4</sup> and R<sup>5</sup> are each independently H, -Cl, -Br, -I, -F or -OC<sub>1-6</sub>alkyl;

J<sup>2</sup> is a member selected from the group consisting of:



Z is  $\text{-NR}^6$ -,  $\text{-O-}$  or  $\text{-S-}$ ;

5  $\text{R}^6$  is H,  $\text{C}_{1-6}$ alkyl or  $\text{C}_{3-8}$ cycloalkyl;


$\text{R}^7$  and  $\text{R}^8$  are independently H,  $\text{-Cl}$ ,  $\text{-Br}$ ,  $\text{-I}$  or  $\text{-F}$ , where at least one of  $\text{R}^7$  and  $\text{R}^8$  is not hydrogen; and

10  $\text{R}^9$  and  $\text{R}^{10}$  are independently H,  $\text{-Cl}$ ,  $\text{-Br}$ ,  $\text{-I}$  or  $\text{-F}$ , where at least one of  $\text{R}^9$  and  $\text{R}^{10}$  is not hydrogen;

and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

15 2. A compound of claim 1, wherein:

A is a member selected from the group consisting of:



20 R is, in each occurrence, independently, H,  $\text{-C}_{1-6}$ alkyl,  $\text{-C}_{3-8}$ cycloalkyl,  $\text{-C}_{0-6}$ alkyl- $\text{OC}_{1-6}$ alkyl,  $\text{-C}_{0-6}$ alkyl- $\text{O}(\text{CH}_2)_{1-4}$ - $\text{COOH}$ ,  $\text{-C}_{0-6}$ alkyl- $\text{O}(\text{CH}_2)_{1-4}$ - $\text{C}(=\text{O})\text{OC}_{1-6}$ alkyl,  $\text{-C}_{0-6}$ alkyl $\text{COOH}$ ,  $\text{-C}_{0-6}$ alkyl $\text{CO}_2\text{C}_{1-6}$ alkyl,  $\text{-C}_{0-6}$ alkyl $\text{OC}_{1-6}$ alkyl,  $\text{-C}_{1-6}$ alkyl $\text{OH}$ ,  $\text{-C}_{0-6}$ alkyl $\text{CONH}_2$ ,  $\text{-C}_{0-6}$ alkyl $\text{CONHC}_{0-6}$ alkyl,  $\text{-C}_{0-6}$ alkyl $\text{CON}(\text{C}_{0-6}$ alkyl) $_2$ ,



-C<sub>0-6</sub>alkylCON(CH<sub>2</sub>)<sub>2-6</sub>, -C<sub>0-6</sub>alkylCON(CH<sub>2</sub>CH<sub>2</sub>)<sub>2</sub>O, -C<sub>0-6</sub>alkylCON(CH<sub>2</sub>CH<sub>2</sub>)<sub>2</sub>SO<sub>2</sub>  
 -C<sub>0-6</sub>alkylCONHaryl, -C<sub>0-6</sub>alkylNH<sub>2</sub>, -C<sub>0-6</sub>alkylNH(C<sub>1-6</sub>alkyl) or  
 -C<sub>0-6</sub>alkylN(C<sub>1-6</sub>alkyl)<sub>2</sub>.


5 R<sup>1</sup> is H, -Cl, -Br, -I or -F, -OCF<sub>3</sub>, -OMe, NH<sub>2</sub>, NHMe, NHMe<sub>2</sub>, -NHCOMe, -  
 NHSO<sub>2</sub>Me; and

R<sup>3</sup> is H, -Cl, -Br, -I, -F, -OC<sub>1-6</sub>alkyl, -NHC<sub>1-6</sub>acyl, -NO<sub>2</sub>, -NHSO<sub>2</sub>C<sub>1-4</sub>alkyl, -CN or  
 -O(CH<sub>2</sub>)<sub>1-4</sub>-COOH.

10

3. A compound of claim 1, wherein:

A is a member selected from the group consisting of:




15 R is, in each occurrence, independently, H, -C<sub>1-6</sub>alkyl, -C<sub>3-8</sub>cycloalkyl, -C<sub>0-6</sub>alkyl-OC<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkyl-O(CH<sub>2</sub>)<sub>1-4</sub>-COOH, -C<sub>0-6</sub>alkyl-O(CH<sub>2</sub>)<sub>1-4</sub>-C(=O)OC<sub>1-6</sub>alkyl,  
 -C<sub>0-6</sub>alkylCOOH, -C<sub>0-6</sub>alkylCO<sub>2</sub>C<sub>1-6</sub>alkyl, -C<sub>0-6</sub>alkylOC<sub>1-6</sub>alkyl, -C<sub>1-6</sub>alkylOH,  
 -C<sub>0-6</sub>alkylCONH<sub>2</sub>, -C<sub>0-6</sub>alkylCONHC<sub>0-6</sub>alkyl, -C<sub>0-6</sub>alkylCON(C<sub>0-6</sub>alkyl)<sub>2</sub>,  
 -C<sub>0-6</sub>alkylCON(CH<sub>2</sub>)<sub>2-6</sub>, -C<sub>0-6</sub>alkylCON(CH<sub>2</sub>CH<sub>2</sub>)<sub>2</sub>O, -C<sub>0-6</sub>alkylCON(CH<sub>2</sub>CH<sub>2</sub>)<sub>2</sub>SO<sub>2</sub>  
 20 -C<sub>0-6</sub>alkylCONHaryl, -C<sub>0-6</sub>alkylNH<sub>2</sub>, -C<sub>0-6</sub>alkylNH(C<sub>1-6</sub>alkyl) or  
 -C<sub>0-6</sub>alkylN(C<sub>1-6</sub>alkyl)<sub>2</sub>;

R<sup>1</sup> is H, -Cl, -Br, -I or -F, -OMe, NH<sub>2</sub>, NHMe, NHMe<sub>2</sub>, -NHCOMe, -NHSO<sub>2</sub>Me;

25 J<sup>1</sup> is a member selected from the group consisting of:

105




X is O or S;

R<sup>3</sup> is H, -Cl, -Br, -I or -F;

5

R<sup>5</sup> is H, -Cl, -Br, -I or -F;

J<sup>2</sup> is a member selected from the group consisting of:



10 Z is -NR<sup>6</sup>-, -O- or -S-;


R<sup>6</sup> is a H, C<sub>1-6</sub> alkyl or C<sub>3-8</sub> cycloalkyl;

R<sup>7</sup> and R<sup>8</sup> are each independently -Cl, -Br, -I or -F; and

15







R<sup>9</sup> and R<sup>10</sup> are each independently -Cl, -Br, -I or -F.

4. A compound of claim 1 of formula (I) having the following structure:







wherein:

20 A is a member selected from the group consisting of:
















5






5

10




5

10




5. A compound of claim 1 of formula (I) having the following structure:








wherein:






A is a member selected from the group consisting of:




110








5








10








5



10








5


6. A compound of claim 1 of formula (I) having the following structure:




10

wherein:


Q is a member selected from the group consisting of:





5


7. A compound of claim 1 of formula (I) having the following structure:








10






wherein:

A is a member selected from the group consisting of:






5



115




5

10


116

5



10


15 8. A compound of claim 1 of formula (I) having the following structure:



wherein:

- 5 R is independently selected from the group consisting of:  
 H, -CO<sub>2</sub>H, -CO<sub>2</sub>Me, -CONH<sub>2</sub>, -CONHMe, -CONHMe<sub>2</sub>, -CON(CH<sub>2</sub>)<sub>4</sub>, -  
 CON(CH<sub>2</sub>)<sub>5</sub>, -CH<sub>2</sub>OH, -CH<sub>2</sub>OMe, -CH<sub>2</sub>CO<sub>2</sub>H, -CH<sub>2</sub>CO<sub>2</sub>Me, -CH<sub>2</sub>CONH<sub>2</sub>,  
 -CH<sub>2</sub>CH<sub>2</sub>OH, -CH<sub>2</sub>CH<sub>2</sub>OMe, -CH<sub>2</sub>NH<sub>2</sub>, -CH<sub>2</sub>N(Me)<sub>2</sub>, and -CH<sub>3</sub>,

- 10 9. A compound of claim 1 of formula (I) having the following structure:




wherein:

R is independently selected from the group consisting of:


- 15 H, -CO<sub>2</sub>H, -CO<sub>2</sub>Me, -CONH<sub>2</sub>, -CONHMe, -CONHMe<sub>2</sub>, -CON(CH<sub>2</sub>)<sub>4</sub>, -  
 CON(CH<sub>2</sub>)<sub>5</sub>, -CH<sub>2</sub>OH, -CH<sub>2</sub>OMe, -CH<sub>2</sub>CO<sub>2</sub>H, -CH<sub>2</sub>CO<sub>2</sub>Me, -CH<sub>2</sub>CONH<sub>2</sub>,  
 -CH<sub>2</sub>CH<sub>2</sub>OH, -CH<sub>2</sub>CH<sub>2</sub>OMe, -CH<sub>2</sub>NH<sub>2</sub>, -CH<sub>2</sub>N(Me)<sub>2</sub>, and -CH<sub>3</sub>,


10. A compound of claim 1 of formula (I) having the following structure:



20 wherein:

A is a member selected from the group consisting of:







5

10






121



11. A compound of claim 1 of formula (I) having the following structure:


5




wherein:

$J^2$  is a member selected from the group consisting of:

10




12. A compound of claim 1 of formula (II) having the following structure:




15


wherein:



















J<sup>1</sup> is a member selected from the group consisting of:





5 13. A compound selected from the group consisting of:






















and all pharmaceutically acceptable isomers, salts, hydrates, solvates and prodrug derivatives thereof.

5

14. A pharmaceutical composition for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound of one of claims 1-13.

10

15. A method for preventing or treating a condition in a mammal characterized by undesired thrombosis comprising administering to said mammal a therapeutically effective amount of a compound of one of claims 1-13.

15 16. The method of claim 15, wherein the condition is selected from the group consisting of:

acute coronary syndrome, myocardial infarction, unstable angina, refractory angina, occlusive coronary thrombus occurring post-thrombolytic therapy or post-coronary angioplasty, a thrombotically mediated cerebrovascular syndrome, embolic  
20 stroke, thrombotic stroke, transient ischemic attacks, venous thrombosis, deep venous thrombosis, pulmonary embolus, coagulopathy, disseminated intravascular coagulation, thrombotic thrombocytopenic purpura, thromboangiitis obliterans, thrombotic disease associated with heparin-induced thrombocytopenia, thrombotic complications associated with extracorporeal circulation, thrombotic complications associated with  
25 instrumentation such as cardiac or other intravascular catheterization, intra-aortic balloon pump, coronary stent or cardiac valve, and conditions requiring the fitting of prosthetic devices.

17. A method for inhibiting the coagulation of biological samples comprising the administration of a compound of one of claims 1-13.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 01/30313

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D409/12 C07D409/14 C07D241/08 C07D403/10 C07D401/14  
C07D491/113 C07D401/12 C07D405/14 A61K31/495 A61P7/02  
/(C07D491/113, 221:00, 317:00)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 40075 A (TAKEDA CHEMICAL INDUSTRIES, LTD.) 12 August 1999 (1999-08-12) claims 1-28 - & EP 1 054 005 A (TAKEDA CHEMICAL INDUSTRIES, LTD.) 22 November 2000 (2000-11-22)	1-17
Y	WO 00 08005 A (MERCK PATENT GMBH) 17 February 2000 (2000-02-17) claims 1-9	1-17
Y	WO 99 37304 A (RHONE-POULENC RORER PHARMACEUTICALS INC.) 29 July 1999 (1999-07-29) cited in the application claims 1-72	1-17
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*G\* document member of the same patent family

Date of the actual completion of the international search

21 February 2002

Date of mailing of the international search report

05/03/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Herz, C

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 01/30313

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, Y	WO 01 07436 A (AVENTIS PHARMACEUTICALS PRODUCTS INC.) 1 February 2001 (2001-02-01) claims 1-31 ----	1-17
Y	WO 00 32590 A (AVENTIS PHARMACEUTICALS PRODUCTS INC.) 8 June 2000 (2000-06-08) cited in the application claims 1-25 ----	1-17
Y	WO 98 21188 A (ZENECA LTD.) 22 May 1998 (1998-05-22) claims 1-18 ----	1-17
A	WO 99 33805 A (MOCHIDA PHARMACEUTICAL CO., LTD.) 8 July 1999 (1999-07-08) claims 1-16 ----	1-17
A	WO 99 06395 A (ZENECA LTD.) 11 February 1999 (1999-02-11) claims 1-9 -----	1-17

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/30313

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9940075	A	12-08-1999	AU 2298899 A	23-08-1999
			CN 1290254 T	04-04-2001
			EP 1054005 A1	22-11-2000
			JP 2000204081 A	25-07-2000
			WO 9940075 A1	12-08-1999
WO 0008005	A	17-02-2000	DE 19835950 A1	10-02-2000
			AU 5411499 A	28-02-2000
			WO 0008005 A2	17-02-2000
WO 9937304	A	29-07-1999	AU 2653399 A	09-08-1999
			BG 104633 A	30-03-2001
			BR 9907300 A	24-10-2000
			CN 1291892 T	18-04-2001
			EE 200000435 A	15-02-2002
			EP 1051176 A1	15-11-2000
			JP 2002501024 T	15-01-2002
			NO 20003808 A	26-09-2000
			PL 342243 A1	04-06-2001
			SK 11082000 A3	18-01-2001
			TR 200002182 T2	21-12-2000
			WO 9937304 A1	29-07-1999
			ZA 9900607 A	27-07-1999
WO 0107436	A	01-02-2001	AU 1923300 A	19-06-2000
			AU 6462800 A	13-02-2001
			WO 0107436 A2	01-02-2001
			WO 0032590 A1	08-06-2000
WO 0032590	A	08-06-2000	AU 1923300 A	19-06-2000
			WO 0032590 A1	08-06-2000
			AU 6462800 A	13-02-2001
			WO 0107436 A2	01-02-2001
			AU 2653399 A	09-08-1999
			BG 104633 A	30-03-2001
			BR 9907300 A	24-10-2000
			EE 200000435 A	15-02-2002
			EP 1051176 A1	15-11-2000
			JP 2002501024 T	15-01-2002
			NO 20003808 A	26-09-2000
			SK 11082000 A3	18-01-2001
WO 9821188	A	22-05-1998	AU 731929 B2	05-04-2001
			AU 4874897 A	03-06-1998
			BG 103430 A	31-07-2000
			BR 9712672 A	26-10-1999
			CN 1235597 A	17-11-1999
			CZ 9901634 A3	11-08-1999
			EP 0937048 A1	25-08-1999
			WO 9821188 A1	22-05-1998
			HU 0001098 A2	28-06-2001
			JP 2001504113 T	27-03-2001
			NO 992230 A	07-05-1999
			PL 333241 A1	22-11-1999
			SK 61399 A3	14-02-2000
			TR 9901006 T2	23-08-1999
			US 6300330 B1	09-10-2001
			ZA 9710062 A	08-05-1998

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/30313

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9933805	A	08-07-1999	AU 1692399 A	19-07-1999
			EP 1048652 A1	02-11-2000
			WO 9933805 A1	08-07-1999
WO 9906395	A	11-02-1999	AU 8547198 A	22-02-1999
			EP 1000057 A1	17-05-2000
			WO 9906395 A1	11-02-1999
			JP 2001512118 T	21-08-2001
			US 6335341 B1	01-01-2002